



Standardised and structured reporting in fetal magnetic resonance imaging: recommendations from the Fetal Task Force of the European Society of Paediatric Radiology

Carmelo Sofia¹ · Michael Aertsen² · Catherine Garel³ · Marie Cassart⁴

Received: 16 May 2024 / Revised: 17 July 2024 / Accepted: 18 July 2024

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2024

Abstract

Over the last decades, magnetic resonance imaging (MRI) has emerged as a valuable adjunct to prenatal ultrasound for evaluating fetal malformations. Several radiological societies advocate for standardised and structured reporting practices to enhance the uniformity of imaging language. Compared to narrative formats, standardised and structured reports offer enhanced content quality, minimise reader variability, have the potential to save reporting time, and streamline the communication between specialists by employing a shared lexicon. Structured reporting holds promise for mitigating medico-legal liability, while also facilitating rigorous scientific data analyses and the development of standardised databases. While structured reporting templates for fetal MRI are already in use in some centres, specific recommendations and/or guidelines from international societies are scarce in the literature. The purpose of this paper is to propose a standardised and structured reporting template for fetal MRI to assist radiologists, particularly those with less experience, in delivering systematic reports. Additionally, the paper aims to offer an overview of the anatomical structures that necessitate reporting and the prevalent normative values for fetal biometrics found in current literature.

Graphical Abstract

Standardised and structured reporting in fetal magnetic resonance imaging: recommendations from the fetal task force of the European Society of Paediatric Radiology

In this paper the authors provide:

1. A standardised template for describing fetal anomalies observed on fetal magnetic resonance imaging
2. An overview of the normative values available in the current literature for various organs and anatomical structures

A standardised and structured reporting template may assist pediatric radiologists in delivering proper systematic reports in fetal magnetic resonance imaging

Pediatric Radiology

Carmelo S., et al. 2024

Keywords Biometry · Fetus · Magnetic resonance imaging · Prenatal · Structured report

Introduction

Over the last decades, magnetic resonance imaging (MRI) has undergone significant technical advancements, rendering it a valuable adjunct to prenatal ultrasound (US) for evaluating fetal malformations. MRI furnishes

supplementary information critical for guiding prenatal counselling and postnatal management decisions.

As per recommendations from the “Fetal Task Force” of the European Society of Paediatric Radiology (ESPR), various indications exist for conducting fetal MRI, encompassing both central nervous system and body pathologies [1, 2]. Numerous medical (radiological and non-radiological) societies recommend the adoption of standardised and

Extended author information available on the last page of the article

structured reporting practices to enhance the consistency and reproducibility of the imaging language [3–14].

While fetal MRI structured report templates are already implemented in some centres, particularly for central nervous system examinations, specific recommendations or guidelines by international societies in the literature are limited [15].

The purpose of this paper is to propose a standardised and structured reporting template in fetal MRI to assist radiologists, particularly those with less experience, in delivering proper systematic reports. In addition to covering general information such as indications, technique, and image quality, we also provide an overview of the anatomical structures that necessitate reporting and the most prevalent fetal biometric data.

Aiming to develop this report template, the authors conducted a survey among the Fetal Task Force members of the ESPR. This questionnaire sought to gather information on the local implementation of structured reporting and the biometric data included in their reports, along with corresponding references from the literature.

Fourteen out of 20 members responded to the questionnaire. The resulting structured report template and the summary of provided fetal biometric data are a concise reflection of the survey's findings and encapsulate the daily reporting practices of these members.

Structured report in fetal magnetic resonance imaging

General information

Indications

A tailored investigation in a proper clinical setting is imperative, and every fetal MRI report should encompass relevant clinical data and family history, especially if related to fetal anomalies. Additionally, informative laboratory test results and, when accessible, genetic data should be included. Conducting a fetal MRI scan following a second-line US scan is mandatory, as it enables a more focused examination and facilitates precise answers to be given to specific questions [1, 2].

The prior ultrasound report should always be available in full text before the MRI examination and the findings prompting a fetal MRI should be summarised in the indications.

This section should also provide information on whether a fetal brain or body MRI (or both) examination is being performed, according to the specific anomalies that have to be clarified. This approach should be discussed with the

referring physician and depends on findings including but not limited to the US scan, maternal history and laboratory results.

Technique

Each report should include technical data: field strength, sequences (also if advanced ones—e.g. diffusion tensor imaging, echoplanar- fluid-attenuated inversion recovery—are performed), sedation if used (drug name and dose).

Image Quality

A visual rating system (low-fair-good–excellent), while subjective in nature, should consistently be provided to assess the image quality and reliability of the examination.

Comparison

When a prior MRI examination is accessible, it should be noted in the text, and a comparison should be conducted to inform the reader of any progression, stability, or regression of previously identified anomalies, as well as the emergence of new ones.

Gestational age

The report should consistently include the gestational age because various fetal growth landmarks are gestational age-dependent, and abnormal findings may suggest fetal growth abnormalities.

Additionally, mentioning the fetal position is crucial as it influences the image quality (for instance a breech presentation complicates brain examination due to maternal respiratory movements).

Fetal life supporting system

Amniotic fluid

A subjective evaluation of the volume of amniotic fluid (normal, increased, reduced, absent) should be noted in the MRI report. Polyhydramnios may result in increased fetal motion, whereas oligohydramnios enhances the value of fetal MRI compared to US.

Placenta

The placental position should be mentioned as well as its heterogeneity, which increases with gestational age.

Fetus

Each section of the report concerns a specific anatomical region, and it should include two parts:

- Description of different structures and anomalies in terms of biometry (when feasible) and morphology. Reference data related to biometry are available in the literature [16–32] and summarised in Table 1 according to the related structure. Free tools and software for comparison of images and percentile calculation are also available online [33, 34].
- Interpretation of imaging findings and conclusion.

Below is an outline of the items that should be checked for each anatomical area, as succinctly summarised in the template (Table 2), along with a brief mention of potential pathologies. This list is necessarily not exhaustive and should be tailored to the pathological context.

Fetal brain and skull

Knowledge of the normal development of the brain is crucial for an accurate report, given the dynamic processes of gyration, cortical maturation, and myelination throughout gestation [35].

All brain structures should be described in terms of presence, appearance (e.g. normal, agenesis, hypoplasia, dysplasia, signal intensity) and biometry.

A systematic evaluation of the following structures is mandatory: the pericerebral spaces, the cortical ribbon, the cerebral parenchyma, the subependymal area, the ventricular

walls, the ventricles, the midline structures, the posterior fossa, the vascular structures.

Pericerebral space: size (subjective evaluation), appearance (possible identification of arachnoid cyst, haematoma, vascular malformations).

Sulcation anomalies: type (gyration delay, thickened cortex, polymicrogyria, abnormal sulci ...), location, extension [36].

Brain parenchyma: appearance and signal intensity (haemorrhage, ischemia, including schizencephaly), volume (subjective evaluation), white matter, mass, basal ganglia.

Subependymal area: appearance (pseudo cysts, haemorrhage, heterotopia).

Lateral ventricles: size, shape regular, irregular (deformation in porencephalic communicating cavity), content (haemorrhage).

Midline structures: anomalies of the corpus callosum in shape and size (partial or complete agenesis, short, thin, thick corpus callosum). *Midline anomalies:* signal intensity (pericallosal lipoma), shape and size of the interhemispheric fissure (incomplete, distorted, displaced, widened) or space-occupying lesion (cyst), septal anomaly, third ventricle (size, shape, position), olfactory bulbs and sulci, optical chiasma, pituitary stalk and gland.

Posterior fossa: amount of pericerebellar fluid (arachnoid cyst, cisterna magna); tentorium (orientation and insertion). **Cerebellum:** size and appearance of the cerebellar hemispheres (haemorrhage, ischemia, dysplasia, mass), the vermis (orientation, partial or complete agenesis, hypoplasia, ischemia), the fourth ventricle (shape, position and content), and the brainstem (shape, bulge of the pons).

Vascular malformations: type, location.

Table 1 Summary of the main fetal biometric reference data reported in the literature

	References: authors (year)	Resources	
Brain	Tilea et al. (2009) [16]	Brain biometry > 26 weeks of gestation	
	Harreld et al. (2011) [17]	Corpus callosum biometry	
	Conte et al. (2018) [18]	Brain biometry > 20 < 24 weeks of gestation	
	Dovjak et al. (2021) [19]	Posterior fossa biometry	
Maxillo-facial region	Robinson et al. (2008) [27]	Orbit and eye biometry	
	Paquette et al. (2009) [20]		
	Nemec et al. (2015) [23] Kooiman et al. (2018) [26]	Mandibular biometry	
Thorax	Rypens et al. (2001) [28] Cannie et al. (2008) [30] Meyers et al. (2018) [25]	Lung volumetry	
	Abdomen	Hyde et al. (2020) [31]	Meconium/large bowel width
		Witzani et al. (2006) [24] Michielsen et al. (2010) [29]	Kidney volumetry
Van Vuuren et al. (2012) [32]		Kidney, adrenals, renal pelvis biometry	
Nemec et al. (2011) [21] Nemec et al. (2012) [22]		Female external genitalia biometry Penile biometry	

Table 2 Fetal magnetic resonance imaging structured report template

Structured fetal magnetic resonance imaging report template

INDICATION**TECHNIQUE****IMAGE QUALITY**
 Low Fair Good Excellent
COMPARISON (date)**FETAL DATA**

- GESTATIONAL AGE
- FETAL POSITION

FETAL LIFE SUPPORTING SYSTEM

- AMNIOTIC FLUID Normal Absent Increased Reduced
- PLACENTA Position (to be detailed)

FETAL ANATOMY**BRAIN**

BRAIN FRONTO-OCCIPITAL DISTANCE mm percentile

SKULL FRONTO-OCCIPITAL DISTANCE mm percentile

CEREBRAL BIPARIETAL DIAMETER mm percentile

BONE BIPARIETAL DIAMETER mm percentile

CORPUS CALLOSUM LENGTH mm percentile

TRANSVERSE DIAMETER OF THE CEREBELLUM mm percentile

VERMIS

- **HEIGHT** mm percentile

- **ANTEROPosterior DIAMETER** mm percentile

- **AREA** mm² percentile

VENTRICLES DIAMETERS (R/L) (if required) mm percentile

Table 2 (continued)

TEGMENTO-VERMIAN ANGLE/ PONS AP DIAMETER (if required)

PERICEREBRAL SPACES	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
SULCATION	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
PARENCHYMA	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
SUBEPENDYMAL AREA	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
LATERAL VENTRICLES	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

MIDLINE STRUCTURES:

- CORPUS CALLOSUM	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- CAVUM SEPTUM PELLUCIDUM	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- OLFACTORY BULBS AND SULCI	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- PITUITARY GLAND	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

POSTERIOR FOSSA:

- BRAINSTEM	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- VERMIS	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- CEREBELLAR HEMISPHERES	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
- POSTERIOR FOSSA FLUID	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

VASCULAR ANOMALIES	<input type="checkbox"/> Absent	<input type="checkbox"/> Present (to be detailed)	<input type="checkbox"/> Cannot be assessed
---------------------------	---------------------------------	---	---

SPINE & SPINAL CORD

SPINAL CORD	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
SPINE	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

MAXILLO-FACIAL REGION

OROFACIAL CLEFT	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
NOSE	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
ORBITS & EYE-BALLS	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

In case of hypo/hypertelorism:

Ocular diameter (OD)

Interocular distances (ID)

Binocular distance (BD)

OPTIC CHIASMA	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed
LIPS	<input type="checkbox"/> Normal	<input type="checkbox"/> Abnormal (to be detailed)	<input type="checkbox"/> Not depicted	<input type="checkbox"/> Cannot be assessed

Table 2 (continued)

MANDIBLE Normal Abnormal (to be detailed) Not depicted Cannot be assessed

MAXILLA Normal Abnormal (to be detailed) Not depicted Cannot be assessed

In case of micrognathia:

Anterior–posterior diameter (APD)

Biparietal diameter (BPD)

Inferior facial angle (IFA)

Jaw index

Oropharyngeal space

Fronto-occipital diameter (FOD)

EARS

EXTERNAL Normal Abnormal (to be detailed) Not depicted Cannot be assessed

MIDDLE Normal Abnormal (to be detailed) Not depicted Cannot be assessed

INNER Normal Abnormal (to be detailed) Not depicted Cannot be assessed

FACIAL MASSES Absent Present (to be detailed) Cannot be assessed

NECK

THYROID Normal Abnormal (to be detailed) Not depicted Cannot be assessed

UPPER AIRWAYS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

In case of atresia or congenital high airway obstruction syndrome (CHAOS):

Gap length (mm)

MASSES Absent Present (to be detailed) Cannot be assessed

THORAX

THYMUS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

ESOPHAGUS (on focused investigations) Normal Abnormal (to be detailed) Not depicted Cannot be assessed

TRACHEA Normal Abnormal (to be detailed) Not depicted Cannot be assessed

In case of atresia or congenital high airway obstruction syndrome (CHAOS):

Gap length (mm)

HEART & GREAT VESSELS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

Table 2 (continued)

LUNGS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

In case of congenital diaphragmatic hernia (CDH) and congenital lung malformations (CLMs):
 Total fetal lung volume (TFLV)
 Expected/observed total fetal lung volume ratio (e/o TFLV)

DIAPHRAGM Normal Abnormal (to be detailed) Not depicted Cannot be assessed

MEDIASTINAL MASSES Absent Present (to be detailed) Cannot be assessed

PLEURAL EFFUSION (Y/N)

PERICARDIAL EFFUSION (Y/N)

ABDOMEN

LIVER Normal Abnormal (to be detailed) Not depicted Cannot be assessed

GALLBLADDER Normal Abnormal (to be detailed) Not depicted Cannot be assessed

SPLEEN Normal Abnormal (to be detailed) Not depicted Cannot be assessed

STOMACH Normal Abnormal (to be detailed) Not depicted Cannot be assessed

SMALL INTESTINE Normal Abnormal (to be detailed) Not depicted Cannot be assessed

Caliber:

COLON Normal Abnormal (to be detailed) Not depicted Cannot be assessed

Caliber:

Meconium 10 mm below the bladder Y/N

ADRENALS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

KIDNEYS Normal Abnormal (to be detailed) Not depicted Cannot be assessed

RENAL PELVIS & Normal Abnormal (to be detailed) Not depicted Cannot be assessed

URETERS

Antero-posterior diameter (APD) of the renal pelvis (R/L)

Ureteral caliber (R/L)

BLADDER Normal Abnormal (to be detailed) Not depicted Cannot be assessed

PELVIC MASSES Absent Present (to be detailed) Cannot be assessed

Table 2 (continued)

ASCITES (Y/N)

ABDOMINAL WALL

 Normal
 Abnormal (to be detailed)
 Not depicted
 Cannot be assessed
SKELETON

EXTREMITIES

- ARMS

 Normal
 Abnormal (to be detailed)
 Not depicted
 Cannot be assessed

- LEGS

 Normal
 Abnormal (to be detailed)
 Not depicted
 Cannot be assessed

OTHER BONES NOT PREVIOUSLY SPECIFIED (if abnormal)

(to be detailed)

CONCLUSIONS

Skull: size (macrocrania, microcephaly) and shape (frontal bossing, cloverleaf skull).

Fetal spinal cord and spine

Although US offers higher spatial resolution, MRI may be valuable for analysing the spinal cord, spinal canal and spine.

Special consideration should be given to the morphology of the *spine* (neural tube defects open/closed type, presence or absence of a sac and its content, spinal canal widening, abnormal curvature, vertebral anomalies, partial agenesis, intracanal mass); presence or absence of a subcutaneous mass including its signal intensity and measurements (lipoma, cyst); and anomalies involving the *spinal cord* (level and appearance of the distal end, possible diastematomyelia) [37].

In case of a *presacral mass*: size, signal intensity (solid/cystic/mixed), location, extension (percentage of intra- or extra-pelvic development), involvement of the spinal canal, impact on the abdominal organs (urinary tract) should be described.

Fetal maxillo-facial region

Any changes in skull integrity and shape should be documented, along with any abnormalities affecting the skull base and facial bones.

The size, shape, position, and content of the maxillo-facial structures should be meticulously assessed, with particular attention to identifying asymmetric paired organs [38].

The following anomalies may be observed:

Orofacial clefts: appearance of the alveolar ridge and the soft tissues (micrognathia, maxillary hypoplasia/cleft, cleft lip, tongue position) and bony palate (cleft). Normal mandibular biometric volumetric data are provided in the literature as a reference for an objective evaluation [23, 26].

Orbits and eyeballs: presence, number, appearance and size of the eyeballs (anophthalmia, microphthalmia, coloboma, cyclopia, hypo/hypertelorism, dacryocystocele, cephaloceles). The chiasma should also be checked (optic nerve agenesis/hypoplasia).

Nose: appearance and size (arrhinia, choanal permeability, presence of the olfactory bulbs).

External ear: presence and appearance (anotia, microtia, external auditory canal atresia or hypoplasia). *Middle and inner ear:* tympanic cavity, cochlea, and semicircular canals.

In case of *facial masses*: size, location, internal architecture and signal intensity (cystic/solid, haemorrhage, homogeneous/inhomogeneous), extension (e.g. intracranial, cervical or thoracic), relationship with the surrounding structures (e.g. upper airway, brain structures) [38].

Fetal neck

The *thyroid gland* should be thoroughly examined, assessing its presence, signal intensity, size (goitre), and position (ectopia).

Upper airways should be identified as non-dilated fluid-filled structures connecting with the lower airways. In case of an interruption, the location and the length of the gap should be assessed.

In cases of *cervical masses*, it is essential to describe the size, signal intensity (whether cystic/solid, homogeneous/inhomogeneous, and presence of fat/blood/calcification components), and extension (into the mediastinum, face, or tongue). Moreover, the airway patency must be evaluated if an ex utero intrapartum treatment procedure is planned.

Fetal thorax

Normal lungs typically exhibit T2 hyperintensity and T1 hypointensity owing to their fluid content. The diaphragm appears as a thin T2 hypointense structure.

In the event of a *congenital diaphragmatic hernia*, it is crucial to identify herniated structures and to report the volume of herniated liver [25, 28, 30, 39, 40].

As part of prognostic evaluation, it is recommended to calculate the total fetal lung volume and expected/observed total fetal lung volume ratio [25, 28, 30].

The *oesophagus* is rarely visible over its entire length on routine examinations. However, it should be specifically sought on dynamic scans centred on the mediastinum in fetuses suspected of having oesophageal atresia, characterised by a blind-ending cervical pouch [41].

Cardiac situs, axis, and size with cardiothoracic index should always be checked and reported if abnormal. If *congenital heart disease* is suspected, the report should detail the cardiac anatomy [42, 43].

The *thymus* is typically situated in the anterior mediastinum. If necessary, such as in cases of agenesis or hypoplasia, its size can be assessed [44].

In case of a *congenital lung malformation/mass*: location, size, morphology, internal structure (cystic, solid, homogeneous, heterogeneous), possible mass effect on the ipsilateral diaphragm and on the mediastinum, pulmonary lobe(s) involved, arterial supply, and venous drainage (e.g. bronchopulmonary sequestration) should be assessed, providing total fetal lung volume and expected/observed total fetal lung volume ratio.

In case of a *mediastinal mass* (e.g. foregut duplication cysts, lymphatic malformations, teratoma, goitre): size, location, extension (e.g. neck, thoracic wall), morphology, characteristics and/or internal structure (cystic/solid, homogeneous/inhomogeneous, fat/blood/calcification components), presence of an infiltrative pattern (in microcystic

lymphatic malformations), compression of surrounding structures (great vessels, trachea, oesophagus, heart, lungs), possible restricted diffusion of solid components on diffusion weighted imaging (DWI), and apparent diffusion coefficient (ADC) maps should be assessed.

Fetal abdomen

Digestive system

The *liver* should be evaluated in terms of size, signal intensity (hypointensity in both T2- and T1-weighted sequences may indicate iron overload), and parenchymal homogeneity (any inhomogeneity may suggest an intraparenchymal mass, which should be described in terms of location, size, morphology, and internal structure) [45]. Normal liver volumetric data are provided in the literature as a reference for an objective evaluation [46].

The fetal *gallbladder* usually appears as a pear-shaped fluid-filled structure with variable signal intensity depending on the gestational age. There are many normal variants regarding its morphology and dimensions [47]. When the gallbladder is absent or if hilar/perihilar cysts are observed, the diagnosis of biliary atresia may be suggested. In such a context, heterotaxia and polysplenia should be searched for [47].

The *stomach* should be seen as a fluid-filled structure in the left hypochondrium; in some settings (congenital diaphragmatic hernia, eventration), its position is important to evaluate [29].

Moreover, MRI facilitates the evaluation of the normal appearance and position of the *intestinal tract*, which should be described based on a combination of T2- and T1-weighted sequences. The meconium should be evaluated in relation to its T1 hyperintense signal and its extension should be assessed (the distal end of the rectum is normally located at least 10 mm below the bladder neck) [48]. The intestinal calibre correlates with the gestational age; the conspicuity of the meconium signal intensity at any part of the bowel increases with time [31]. Reference ranges of bowel width can be found in Hyde et al. [31]. In cases of gastrointestinal obstruction, it is essential to report the location and signal intensity of the contents of dilated loops and the presence of a micro-colon, as this enables assessment of the level of obstruction [49].

Urogenital system

Kidneys: size, location (ectopia), and parenchymal appearance should be evaluated. In the event of a renal mass, it is essential to describe its size, internal structure, and relationship with the renal hilum. DWI with ADC maps can provide

valuable information about the mass, including any signal restriction indicative of malignancy, as well as the presence of residual functional renal parenchyma [24]. Furthermore, the renal cavities should be analysed, and any anomalies should be reported in cases of suspected uro/nephropathies with pelvi-caliceal and ureteral diameters.

Renal parenchymal evaluation can be conducted by assessing a ratio to the liver or renal pelvis' signal intensity for maturation [24]. ADC maps can help in the identification of functional renal parenchyma [24].

The normal *adrenal gland* is typically visible from around 20 weeks gestational age and tends to be relatively large. The report should state the presence as well as the normal size of the adrenal glands [50]. In cases of suprarenal/adrenal masses: size, appearance (cystic, solid, fatty, homogeneous, heterogeneous, haemorrhagic/proteinaceous contents), DWI restriction—indicative of malignancy, may assist in establishing a specific diagnosis (haemorrhage, sequestration, neuroblastoma) [51].

The *bladder*: presence, size, wall (thick, smooth, or irregular), and contents should be assessed.

While *penile abnormalities* are typically better detected with US, MRI reference ranges for the total penile and outer penile length have been published [22].

In case of a *pelvic mass*, it is important to describe its size, location, extent, contents' signal intensity (such as haemorrhagic, pure fluid, or meconial), and its relationships with the surrounding structures (such as spine, digestive tract, bladder), which may help to characterise the malformation (anorectal malformation, urogenital sinus, or cloacal malformation) or tumour.

Abdominal wall defect (gastroschisis, omphalocele): region of involvement, herniated organs, presence or absence of a peritoneal-amniotic membrane, cord insertion and integrity, and bladder location (e.g. exstrophy) should be reported [52].

Fetal skeleton

While indications for imaging the fetal skeleton with MRI are limited and controversial (with US remaining the gold standard technique), evaluation should be considered within the context of conditions such as spina bifida and complex fetal anomalies. MRI can potentially assess deformities and the alignment of joints or bones (e.g. club foot, scoliosis, craniosynostosis, joint dislocation) [53].

The following anomalies may be encountered: limb shortening, absence (location), positional deformities (location, type), skeletal dysplasia, skull (see the “[Fetal brain and skull](#)” section), and vertebral anomalies (see the “[Fetal spinal cord and spine](#)” section).

Conclusion

The report should end with a concise summary of key findings, including a diagnostic hypothesis and potential syndromes that could guide genetic testing if it has not yet been conducted. Recommendations for follow-up should also be provided, if necessary.

In cases where the examination does not yield conclusive results, this should be clearly stated to facilitate appropriate diagnostic management (e.g. strict imaging follow-up with US and/or repeated MRI, or fetal low-dose CT for skeletal dysplasia) [54].

Discussion

The widespread adoption of structured reporting is essential for delivering the highest quality of service to referring physicians and, ultimately, to patients [9].

Compared to narrative format texts, standardised and structured reports appear to enhance content quality, diminish reader variability, and, through a shared lexicon, facilitate communication among specialists [12, 55, 56]. Each of these elements contributes independently to the more efficient integration of the radiological report into the clinical pathway.

Structured reporting holds the potential to mitigate medico-legal liability and offers advantages for scientific data analysis and the establishment of standardised databases [12, 56].

Structured reports appear to streamline reporting processes and, while challenging to quantify, the enhanced communication facilitated by a well-constructed radiological report can potentially reduce the overall assessment time. This efficiency is attributed to the use of uniform terminology and the inclusion of all necessary information, thereby minimising the need for additional discussions and second readings.

Although some centres share their fetal MRI reporting templates online, these are typically tailored to local experience and practice (<https://www.pedrad.org/Specialties/Fetal-Imaging/Fetal-MRI-General-Information#49043614-templates>). The International Society of Ultrasound in Obstetrics & Gynecology (ISUOG) practice guidelines on fetal MRI provide some general recommendations on what a fetal MRI report should include [15]. However, a comprehensive framework for analysing the fetus using fetal MRI—covering the reporting of both normal and abnormal findings along with organ-specific biometry values—is still absent in the current literature.

More recently, Thater et al. provided a structured report template specifically designed for fetuses with congenital diaphragmatic hernia [57]. However, fetal MRI is now being

applied more broadly to address developmental questions across various organ systems, including facial, thoracic, and gastrointestinal abnormalities, in addition to its established role as a complementary tool for assessing central nervous system abnormalities.

Given the growing significance of fetal MRI and its increasing accessibility [58], our aim is to bridge the current gap by offering a comprehensive framework. This framework is intended to support radiologists who are new to fetal MRI, as well as those with more experience, allowing them to evaluate and refine their current practices. Our proposed template provides a structured approach for the evaluation of all fetal organ systems.

This report template, including the biometric reference data, should be viewed as an initial step. Future validation of this template will be necessary, recognising that the adoption of a new reporting format will require adjustments from both radiologists and referring clinicians.

Conclusion

With this paper, we have provided a standardised template for describing fetal anomalies observed using fetal MRI, along with an overview of the normative values available in the current literature for various organs and anatomical structures. This template may be adjusted according to local procedures and preferences.

Acknowledgements The authors extend their gratitude to the members of the Fetal Task Force of the European Society of Paediatric Radiology for their invaluable participation in the internal survey regarding the use of structured reporting in their daily practice. We acknowledge their contributions as follows: Eleonore Blondiaux, Pablo Caro, Céline Habre, Mai Lan Ho, Raimund Kottke, Martin Kyncl, Georgia Papaioannou, Monica Rebollo, Andrea Rossi, and Elspeth H Whitby.

Author contribution C.S. conceived the study. C.S. and M.A. conducted the literature search and analysed the data. C.S. and M.A. drafted the initial manuscript. C.G. and M.C. reviewed the initial manuscript. All authors reviewed and approved the final manuscript.

Data availability Data sharing is not applicable.

Declarations

Conflicts of interest None

References

- Papaioannou G, Caro-Domínguez P, Klein WM et al (2022) Indications for magnetic resonance imaging of the fetal body (extra-central nervous system): recommendations from the European Society of Paediatric Radiology Fetal Task Force. *Pediatr Radiol*. <https://doi.org/10.1007/s00247-022-05495-4>
- Papaioannou G, Klein W, Cassart M, Garel C (2021) Indications for magnetic resonance imaging of the fetal central nervous system: recommendations from the European Society of Paediatric Radiology Fetal Task Force. *Pediatr Radiol* 51:2105–2114
- Beets-Tan RGH, Lambregts DMJ, Maas M et al (2018) Magnetic resonance imaging for clinical management of rectal cancer: updated recommendations from the 2016 European Society of Gastrointestinal and Abdominal Radiology (ESGAR) consensus meeting. *Eur Radiol* 28:1465–1475
- Andreotti RF, Timmerman D, Strachowski LM et al (2020) O-RADS US Risk Stratification and Management System: a consensus guideline from the ACR Ovarian-Adnexal Reporting and Data System Committee. *Radiology* 294:168–185
- Alfred Witjes J, Max Bruins H, Carrión A et al (2024) European Association of Urology guidelines on muscle-invasive and metastatic bladder cancer: summary of the 2023 guidelines. *Eur Urol* 85:17–31
- Marrero JA, Kulik LM, Sirlin CB et al (2018) Diagnosis, staging, and management of hepatocellular carcinoma: 2018 practice guidance by the American Association for the Study of Liver Diseases. *Hepatol Baltim Md* 68:723–750
- KSAR Study Group for Rectal Cancer (2017) Essential items for structured reporting of rectal cancer MRI: 2016 consensus recommendation from the Korean Society of Abdominal Radiology. *Korean J Radiol* 18:132–151
- Al-Hawary MM, Francis IR, Chari ST et al (2014) Pancreatic ductal adenocarcinoma radiology reporting template: consensus statement of the Society of Abdominal Radiology and the American Pancreatic Association. *Radiology* 270:248–260
- European Society of Radiology (ESR) (2018) ESR paper on structured reporting in radiology. *Insights Imaging* 9:1–7
- Sardanelli F, Fallenberg EM, Clauser P et al (2017) Mammography: an update of the EUSOBI recommendations on information for women. *Insights Imaging* 8:11–18
- Mottet N, van den Bergh RCN, Briers E et al (2021) EAU-EANM-ESTRO-ESUR-SIOG guidelines on prostate cancer-2020 update. Part 1: screening, diagnosis, and local treatment with curative intent. *Eur Urol* 79:243–262
- European Society of Radiology (ESR) (2023) ESR paper on structured reporting in radiology-update 2023. *Insights Imaging* 14:199
- American College of Radiology Committee on LI-RADS®. Liver-RADS Assessment Categories 2018. Available at <https://www.acr.org/-/media/ACR/Files/RADS/LI-RADS/LI-RADS-2018-Core.pdf>. Accessed 3 Jul 2024
- D’Orsi CJ, Sickles EA, Mendelson EB, Morris EA, et al. ACR BI-RADS® Atlas, Breast Imaging Reporting and Data System. Reston, VA, American College of Radiology; 2013.
- Prayer D, Malinger G, De Catte L et al (2023) ISUOG practice guidelines (updated): performance of fetal magnetic resonance imaging. *Ultrasound Obstet Gynecol* 61:278–287
- Tilea B, Alberti C, Adamsbaum C et al (2009) Cerebral biometry in fetal magnetic resonance imaging: new reference data. *Ultrasound Obstet Gynecol* 33:173–181
- Harreld JH, Bhore R, Chason DP, Twickler DM (2011) Corpus callosum length by gestational age as evaluated by fetal MR imaging. *AJNR Am J Neuroradiol* 32:490–494
- Conte G, Milani S, Palumbo G et al (2018) Prenatal brain MR imaging: reference linear biometric centiles between 20 and 24 gestational weeks. *AJNR Am J Neuroradiol* 39:963–967
- Dovjak GO, Schmidbauer V, Brugger PC et al (2021) Normal human brainstem development in vivo: a quantitative fetal MRI study. *Ultrasound Obstet Gynecol* 58:254–263
- Paquette LB, Jackson HA, Tavaré CJ et al (2009) In utero eye development documented by fetal MR imaging. *AJNR Am J Neuroradiol* 30:1787–1791

21. Nemeč SF, Nemeč U, Weber M et al (2011) Female external genitalia on fetal magnetic resonance imaging. *Ultrasound Obstet Gynecol* 38:695–700
22. Nemeč SF, Nemeč U, Weber M et al (2012) Penile biometry on prenatal magnetic resonance imaging. *Ultrasound Obstet Gynecol* 39:330–335
23. Nemeč U, Nemeč SF, Brugger PC et al (2015) Normal mandibular growth and diagnosis of micrognathia at prenatal MRI. *Prenat Diagn* 35:108–116
24. Witzani L, Brugger PC, Hörmann M et al (2006) Normal renal development investigated with fetal MRI. *Eur J Radiol* 57:294–302
25. Meyers ML, Garcia JR, Blough KL et al (2018) Fetal lung volumes by MRI: normal weekly values from 18 through 38 weeks' gestation. *AJR Am J Roentgenol* 211:432–438
26. Kooiman TD, Calabrese CE, Didier R et al (2018) Micrognathia and oropharyngeal space in patients with robin sequence: prenatal MRI measurements. *J Oral Maxillofac Surg* 76:408–415
27. Robinson AJ, Blaser S, Toi A et al (2008) MRI of the fetal eyes: morphologic and biometric assessment for abnormal development with ultrasonographic and clinicopathologic correlation. *Pediatr Radiol* 38:971–981
28. Rypens F, Metens T, Rocourt N et al (2001) Fetal lung volume: estimation at MR imaging-initial results. *Radiology* 219:236–241
29. Michielsen K, Meersschaert J, De Keyzer F et al (2010) MR volumetry of the normal fetal kidney: reference values. *Prenat Diagn* 30:1044–1048
30. Cannie MM, Jani JC, Van Kerkhove F et al (2008) Fetal body volume at MR imaging to quantify total fetal lung volume: normal ranges. *Radiology* 247:197–203
31. Hyde G, Fry A, Raghavan A, Whitby E (2020) Biometric analysis of the foetal meconium pattern using T1 weighted 2D gradient echo MRI. *BJR Open* 2:20200032
32. van Vuuren SH, HaM Damen-Elias, Stigter RH et al (2012) Size and volume charts of fetal kidney, renal pelvis and adrenal gland. *Ultrasound Obstet Gynecol* 40:659–664
33. Fetal Brain Atlas. http://crl.med.harvard.edu/research/fetal_brain_atlas/. Accessed 3 Apr 2024
34. Kyriakopoulou V, Vatansever D, Davidson A et al (2017) Normative biometry of the fetal brain using magnetic resonance imaging. *Brain Struct Funct* 222:2295–2307
35. Glenn OA, Barkovich AJ (2006) Magnetic resonance imaging of the fetal brain and spine: an increasingly important tool in prenatal diagnosis, part 1. *AJNR Am J Neuroradiol* 27:1604–1611
36. Righini A, Genovese M, Parazzini C et al (2020) Cortical formation abnormalities on foetal MR imaging: a proposed classification system trialled on 356 cases from Italian and UK centres. *Eur Radiol* 30:5250–5260
37. Garel J, Rossi A, Blondiaux E et al (2024) Prenatal imaging of the normal and abnormal spinal cord: recommendations from the Fetal Task Force of the European Society of Paediatric Radiology (ESPR) and the European Society of Neuroradiology (ESNR) Pediatric Neuroradiology Committee. *Pediatr Radiol* 54:548–561
38. Nagarajan M, Sharbidre KG, Bhabad SH, Byrd SE (2018) MR imaging of the fetal face: comprehensive review. *Radiographics* 38:962–980
39. Kitano Y, Okuyama H, Saito M et al (2011) Re-evaluation of stomach position as a simple prognostic factor in fetal left congenital diaphragmatic hernia: a multicenter survey in Japan. *Ultrasound Obstet Gynecol* 37:277–282
40. Lazar DA, Ruano R, Cass DL et al (2012) Defining "liver-up": does the volume of liver herniation predict outcome for fetuses with isolated left-sided congenital diaphragmatic hernia? *J Pediatr Surg* 47:1058–1062
41. Salomon LJ, Sonigo P, Ou P et al (2009) Real-time fetal magnetic resonance imaging for the dynamic visualization of the pouch in esophageal atresia. *Ultrasound Obstet Gynecol* 34:471–474
42. Faruk Topaloğlu Ö, Koplay M, Kılınçer A et al (2023) Quantitative measurements and morphological evaluation of fetal cardiovascular structures with fetal cardiac MRI. *Eur J Radiol* 163:110828
43. Saleem SN (2008) Feasibility of MRI of the fetal heart with balanced steady-state free precession sequence along fetal body and cardiac planes. *AJR Am J Roentgenol* 191:1208–1215
44. Story L, Zhang T, Uus A et al (2021) Antenatal thymus volumes in fetuses that delivered <32 weeks' gestation: an MRI pilot study. *Acta Obstet Gynecol Scand* 100:1040–1050
45. Cassart M, Avni FE, Guibaud L et al (2011) Fetal liver iron overload: the role of MR imaging. *Eur Radiol* 21:295–300
46. Hawkins-Villarreal A, Moreno-Espinosa AL, Martinez-Portilla RJ et al (2022) Fetal liver volume assessment using magnetic resonance imaging in fetuses with cytomegalovirus infection†. *Front Med* 9:889976
47. Avni FE, Garel C, Naccarella N, Franchi-Abella S (2023) Anomalies of the fetal gallbladder: pre-and postnatal correlations. *Pediatr Radiol* 53:602–609
48. Saguintaah M, Couture A, Veyrac C et al (2002) MRI of the fetal gastrointestinal tract. *Pediatr Radiol* 32:395–404
49. Furey EA, Bailey AA, Twickler DM (2016) Fetal MR imaging of gastrointestinal abnormalities. *Radiogr Rev Publ Radiol Soc N Am Inc* 36:904–917
50. Smitthimedhin A, Rubio EI, Blask AR et al (2020) Normal size of the fetal adrenal gland on prenatal magnetic resonance imaging. *Pediatr Radiol* 50:840–847
51. Flanagan SM, Rubesova E, Jaramillo D, Barth RA (2016) Fetal suprarenal masses—assessing the complementary role of magnetic resonance and ultrasound for diagnosis. *Pediatr Radiol* 46:246–254
52. Victoria T, Andronikou S, Bowen D et al (2018) Fetal anterior abdominal wall defects: prenatal imaging by magnetic resonance imaging. *Pediatr Radiol* 48:499–512
53. Chauvin NA, Victoria T, Khwaja A et al (2020) Magnetic resonance imaging of the fetal musculoskeletal system. *Pediatr Radiol* 50:2009–2027
54. Bach P, Cassart M, Chami M et al (2023) Exploration of the fetal skeleton by ultra-low-dose computed tomography: guidelines from the Fetal Imaging Task Force of the European Society of Paediatric Radiology. *Pediatr Radiol* 53:621–631
55. Schwartz LH, Panicek DM, Berk AR et al (2011) Improving communication of diagnostic radiology findings through structured reporting. *Radiology* 260:174–181
56. Nobel JM, Kok EM, Robben SGF (2020) Redefining the structure of structured reporting in radiology. *Insights Imaging* 11:10
57. Thater G, Weidner A, Rafat N et al (2024) Structured reporting in fetal magnetic resonance imaging with congenital diaphragmatic hernia. *Prenat Diagn*. <https://doi.org/10.1002/pd.6593>
58. Pesacreta LD, Cilli KE, Lawrence AK, Bulas DI (2020) The key role of the pediatric radiologist in developing a multidisciplinary fetal center. *Pediatr Radiol* 50:1801–1809

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Carmelo Sofia¹  · Michael Aertsen²  · Catherine Garel³  · Marie Cassart⁴ 

✉ Carmelo Sofia
carm.sofia@tiscali.it

¹ Department of Biomedical Sciences and Morphologic and Functional Imaging, University of Messina, Policlinico “G. Martino”, Via Consolare Valeria 1, 98100 Messina, Italy

² Department of Radiology, University Hospitals Katholieke Universiteit (KU), Louvain, Belgium

³ Department of Radiology, Armand-Trousseau Hospital, APHP, Sorbonne University, Paris, France

⁴ Department of Radiology and Fetal Medicine, Iris South Hospitals, Brussels, Belgium