



# Consensus guidelines on the bedside assistant skills required in robotic surgery

Riley Brian<sup>1</sup> · Alyssa Murillo<sup>1</sup> · Camilla Gomes<sup>1</sup> · Hueylan Chern<sup>1</sup> · Daniel Oh<sup>2,3</sup> · Patricia S. O'Sullivan<sup>1</sup>

Received: 21 May 2024 / Accepted: 18 August 2024  
© The Author(s) 2024

## Abstract

**Background** While bedside assistants play a critical role in many robotic operations, substantial heterogeneity remains in bedside assistant training pathways. As such, this study aimed to develop consensus guidelines for bedside assistant skills required for team members in robotic operations.

**Methods** We designed a study using the Delphi process to develop consensus guidelines around bedside assistant skills. We generated an initial list of bedside assistant skills from the literature, training materials, and expert input. We selected experts for the Delphi process based on prior scholarship in the area of robotic bedside assistant education and experience facilitating robotic bedside assistant training. For each item, respondents specified which robotic team members should have the skill from a list of “basic” bedside assistants, “advanced” bedside assistants, surgeons, surgical technologists, and circulating nurses. We conducted two rounds of the Delphi process and defined 80% agreement as sufficient for consensus.

**Results** Fourteen experts participated in two rounds of the Delphi process. By the end of the second round, the group had reached consensus on 253 of 305 items (83%). The group determined that “basic” bedside assistants should have 52 skills and that “advanced” bedside assistants should have 60 skills. The group also determined that surgeons should have 54 skills, surgical technologists should have 25 skills, and circulating nurses should have 17 skills. Experts agreed that all participants should have certain communication skills and basic knowledge of aspects of the robotic system.

**Conclusions** We developed consensus guidelines on the skills required during robotic surgery by bedside assistants and other team members using the Delphi process. These findings can be used to design training around bedside assistant skills and assess team members to ensure that each team member has the appropriate skills. Hospitals can also use these guidelines to standardize expectations for robotic team members.

**Keywords** Robotic surgery · Bedside assistance · Technical skills · Non-technical skills

Bedside assistants play a critical role in many robotic operations [1]. By managing the robotic system and providing adjunct laparoscopic support, bedside assistants can facilitate case progression and patient safety [2]. Multiple studies have suggested that the experience of the bedside assistant affects important outcomes in robotic surgery [3–6]. Thus, education and training pathways for

bedside assistants must adequately prepare them to be effective in the operating room [7].

Despite the importance of appropriate instruction for bedside assistants, existing training programs, case requirements, and credentialing processes vary substantially [7–10]. Many authors report curricula requiring learners to participate in as few as five cases before being considered competent; however, prior learning curve analyses suggest that bedside skills continue to mature over dozens of cases [1]. Structured education in robotic surgery has tended to de-emphasize bedside assistant skills in favor of a greater focus on the training of the console surgeon [11, 12].

Importantly, bedside assistants come from a wide variety of educational backgrounds—including board-certified surgeons, physician assistants, nurses, surgical technologists, and residents [1]. More recently, medical students have

✉ Riley Brian  
riley.brian@ucsf.edu

<sup>1</sup> Department of Surgery, University of California San Francisco, 513 Parnassus Avenue, S-321, San Francisco, CA 94143, USA

<sup>2</sup> Department of Surgery, University of Southern California, Los Angeles, CA, USA

<sup>3</sup> Intuitive Surgical, Sunnyvale, CA, USA

trained to work as bedside assistants [13–15]. In the setting of such diverse expertise, several authors have identified key components of bedside assistant training [16, 17]. Building on these suggestions, both the Association of Surgical Technologists and the Association of Surgical Assistants have published guidelines outlining necessary skills for those working at the robotic bedside [18, 19]. Together, these publications indicate skills needed for bedside assistants in the pre-operative, intra-operative, and post-operative settings [16–19]. Nonetheless, skills described in prior publications vary in scope, highlight different skills, and arose from authors' perceptions.

Prior work has specifically called for standardization of robotic surgical curricula both for education and credentialing purposes [20]. By better defining the skills needed of bedside assistants, we can clarify training pathways and ground credentialing decisions in evidence. As such, this study aimed to develop consensus guidelines from diverse experts for the bedside assistant skills required in robotic operations by specific roles and across educational backgrounds. We hypothesized that experts would identify required skills for bedside assistants and other robotic surgery team members that vary based on role.


## Methods

### Study design

We developed consensus guidelines for the bedside assistant skills required for robotic operations using the Delphi process (Fig. 1) [21]. We chose to use a consensus group method of guideline generation given the lack of trial-based data to support a standardized list, and we selected the Delphi process for reasons of cost and due to the geographic separation of experts [22].

### Item development

We developed an initial list of bedside assistant skills based on prior peer-reviewed and society-based publications suggesting skills useful for bedside assistants [16–19]. We collated all previously published items and adapted them if needed to create consistency of phrasing and eliminate redundancies. Multiple authors attended bedside assistant training courses run by the University of California San Francisco (UCSF) and Intuitive Surgical and discussed the candidate items with external collaborators (see acknowledgements) to identify additional items (Intuitive Surgical, Sunnyvale, CA). For each item, we asked respondents to specify which robotic team members should have the skill from a list of “basic” bedside assistants, “advanced” bedside assistants, surgeons, surgical



**Fig. 1** Evolution of items during the study

technologists, and circulating nurses. We defined “basic” bedside assistants as those working in cases without an assist port and “advanced” bedside assistants as those working in cases with an assist port. We chose to identify skills for non-bedside assistant team members (i.e., surgeons, surgical technologists, and circulating nurses) to allow for comparisons in skill requirements among roles. We used a scale from one to four for each item, which ranged from “definitely does not need the skill” for one to

“definitely needs the skill” for four. Four authors piloted the questionnaire and we made additional modifications prior to finalizing the items (available as Supplementary File 1).

### Expert identification and recruitment

We selected experts based on two major criteria: prior scholarship in the area of robotic bedside assistant education and prior experience conducting robotic bedside assistant training. We identified experts meeting the first criterion through a literature review, and experts meeting the second criterion through discussion with the Intuitive Surgical professional education team. We aimed to recruit a diverse group of experts representing bedside assistants and surgeons from multiple surgical specialties.

After generating an initial list of experts, we individually emailed each candidate and explained the reason for selection and the study’s design and purpose. We offered experts a gift card incentive for each round in which they participated.

### Delphi process

After candidates expressed an interest in participating, we individually disseminated the first round of items with an email containing a Qualtrics link (Qualtrics, Provo, UT). We did not ask participants for the rationale behind their responses. After experts participated in the first round, we collated responses to identify the mean score assigned for each item. Prior to starting the study, we determined that 80% consensus would be required for an item to be included as a skill. We defined 80% consensus as having 80% of respondents state that a team member “probably” or “definitely” needs the skill described by the item based on the four-point scale (Supplementary File 1). We calculated Cronbach’s alpha in each round using StataNow/BE 18.5 for Mac (StataCorp, College Station, TX). We maintained total anonymity among participants throughout the process. Participants did not meet at any time.

We included those items with less than 80% consensus after the first round in the second round. We also incorporated new items based on experts’ suggestions. We then asked the same experts to participate in a second round. If an item did not achieve 80% consensus by the end of the second round, we deigned to not include them item in the final list due to lack of expert clarity about the necessity of the item. We disseminated final results to all participants at the end of the second round.

### Ethical considerations

Our Institutional Review Board determined this study to be exempt from review (IRB23-39,876).

## Results

We recruited 14 experts to participate in the two rounds of the Delphi process. All 14 experts (100%) who expressed interest in participating before starting the process subsequently completed both rounds. Experts included six surgeons, four physician assistants, two certified surgical assistants, one nurse, and one surgical technologist who also worked as a robotic coordinator. Thirteen experts practiced in eight geographically diverse states in the United States while one expert practiced internationally. Experts represented specialties within general surgery, obstetrics-gynecology, and urology. Experts had participated in a median of 1650 robotic cases (IQR 1000-3500).

The review of items resulted in 59 items for consideration across five different team members, resulting in 295 decisions for each expert. All 14 experts completed all 295 selections. In the first round, the group reached consensus on 220 of 295 decisions (75%) with Cronbach’s alpha of 0.98. More specifically, the group reached consensus regarding 47 of 59 items (80%) for “basic” bedside assistant skills and 55 of 59 items (93%) for “advanced” bedside assistant skills. They also reached consensus on 53 of 59 items (89%) for surgeon skills, 21 of 59 items (36%) for surgical technologist skills, and 44 of 59 items (75%) for circulating nurse skills. Experts added two unique skills—related to manually releasing instruments and loading/unloading staplers—to the 59 original skills during the first round of the Delphi process resulting in a list of 61 skills.

In the second round, all 14 experts made 84 of the 85 selections (98.8%). One expert did not respond to one of the items, though the item reached consensus and this would not have changed based on the fourteenth expert’s response. The group reached consensus on 33 of 85 unique items (39%) with Cronbach’s alpha of 0.95. The group reached consensus on 5 of 14 items (36%) for “basic” bedside assistant skills and 5 of 6 items (83%) for “advanced” bedside assistant skills. They also reached consensus on 2 of 8 items (25%) for surgeon skills, 15 of 40 items (38%) for surgical technologist skills, and 6 of 17 items (35%) for circulating nurse skills.

Thus, by the end of the second round, the group had reached consensus on 253 of the 305 unique items (83%). The group determined that “basic” bedside assistants should have 52 skills and that “advanced” bedside assistants should have 60 skills. They also determined that surgeons should have 54 skills, surgical technologists should have 25 skills, and circulating nurses should have 17 skills. Skills spanned pre- and post-operative technical skills (Table 1), intra-operative technical skills (Table 2), and non-technical skills (Table 3). There was substantial

**Table 1** Pre- and post-operative technical skill list (Color table online)

Skill	“Basic” bedside assistant (cases without assist port)	“Advanced” bedside assistant (cases with assist port)	Surgeon	Surgical technologist	Circulating nurse
Connect each component of the da Vinci system (i.e., Patient Cart, Vision Cart, Surgeon Console) to each relevant component	No Consensus	Round Two	No Consensus	Round Two	Round One
Power on and off the da Vinci system	No Consensus	Round One	No Consensus	Round One	Round One
Identify all buttons on each component of the da Vinci system (i.e., Patient Cart, Vision Cart, Surgeon Console)	No Consensus	Round One	Round One	No Consensus	Round One
Place sterile drapes on the da Vinci Patient Cart	Round Two	Round Two	Round Two	Round One	Round Two
Position the patient appropriately for the specific case to avoid collisions or pressure on the patient’s body	Round One	Round One	Round One	Round One	No Consensus
Choose the appropriate settings on the da Vinci Patient Cart for the procedure	No Consensus	Round One	Round One	No Consensus	Round One
Check patient for any tissue trauma from positioning, trocars, or robotic arm interactions	Round One	Round One	Round One	No Consensus	Round Two

In round one and round two of the Delphi process, experts came to consensus on pre- and post-operative technical skills that bedside assistants and other team members should have (green) or do not need (red) in a robotic operation. Experts did not come to consensus on the necessity of team members having a number of the skills (gray)

overlap between skill requirements for bedside assistants and surgeons, with less overlap in skill requirements among bedside assistants, surgical technologists, and circulating nurses.

## Discussion

In this study, we used the Delphi process to develop consensus guidelines on the bedside assistant skills required during robotic surgery. We identified 61 skills related to robotic bedside assistance and experts determined whether bedside assistants and other team members should have each skill. Experts deemed all 61 skills to be required of at least one team member in an operation, though there was no agreement as to which team members needed some of the skills. Overall, these findings can be used to design training around bedside assistant skills and assess team members to ensure that each team member has the appropriate skills. We anticipate that hospitals can use these guidelines—rather than team members’ degrees—to standardize expectations for robotic team members around bedside assistant skills.

We noted several interesting findings with regard to consensus items. Experts agreed that “basic” bedside assistants and “advanced” bedside assistants should have most of the intra-operative technical skills included in the list as candidate items. We speculate that all intra-operative technical skills may be viewed as essential given the possibility of unexpected events during the course of any operation. Experts most strongly agreed on the importance of communication in robotic operations. Given the unique physical set up of the robotic operating room, this finding supports

prior work highlighting the centrality of communication to the success of robotic surgery [23, 24]. Experts disagreed more substantially around whether bedside assistants needed certain pre- and post-operative technical skills and non-technical skills related to knowledge around procedures and patients. Finally, we found quite varied opinions as to the skills required of surgical technologists. This likely reflects the diverse practice settings of the experts included in this study, as well as more general variability in surgical technologists’ role in different procedures. We had chosen to include surgeons, surgical technologists, and circulating nurses as comparator groups to contrast their necessary skills with those of bedside assistants.

This study builds on prior work establishing the necessary skills for bedside assistants and other team members in robotic surgery. One prior consensus study identified possible credentialing requirements in robotic surgery, though limited proposals to operating surgeons [12]. Other guidelines have focused on bedside assistant skills, though these stemmed from authors’ perceptions rather than expert consensus [16–19]. These guidelines formed the basis for many of the items we evaluated in this study. Our work adds to these prior publications by its consensus-based approach to defining necessary skills for bedside assistants. Of note, we did not ask experts to stratify skills for bedside assistants beyond “basic” and “advanced” cases. Certainly, those in different roles (e.g., physician assistant vs surgical resident) and with different experience levels may be expected to demonstrate “basic” or “advanced” skills at various times in their career or training depending on institution-specific case volumes and policies. Institutions may use the guidelines developed here to determine when bedside assistants

**Table 2** Intra-operative technical skill list (Color table online)

Skill	“Basic” bedside assistant (cases without assist port)	“Advanced” bedside assistant (cases with assist port)	Surgeon	Surgical technologist	Circulating nurse
Insert ports	Round Two	Round One	Round One	Round One	Round One
Space ports appropriately to avoid collisions	Round Two	Round One	Round One	No Consensus	Round One
Activate CO2 insufflation and set to appropriate pressure and flow	Round One	Round One	Round One	Round Two	Round One
Set up CO2 humidification	Round One	Round One	No Consensus	Round One	Round One
Set up AirSeal®	Round One	Round One	Round One	Round One	Round One
Set up smoke evacuation system	Round One	Round One	No Consensus	Round One	Round Two
Drive the da Vinci robot to the patient and adjust boom positioning	No Consensus	Round One	Round One	No Consensus	Round One
Perform targeting to the appropriate anatomy	Round One	Round One	Round One	No Consensus	Round One
Dock the robotic arms to each trocar	Round One	Round One	Round One	Round One	Round One
“Burp” each trocar using the port clutch	Round One	Round One	Round One	No Consensus	Round One
Optimize clearance of arms and joints	Round One	Round One	Round One	No Consensus	Round One
Optimize alignment of the camera port, target anatomy, and center column to prevent collisions	Round One	Round One	Round One	No Consensus	Round One
Attach energy sources to instruments	Round One	Round One	Round One	Round One	No Consensus
Adjust energy settings	Round One	Round One	Round One	No Consensus	Round One
Manipulate the robotic arms using the port and instrument clutches	Round One	Round One	Round One	No Consensus	Round One
Load and insert robotic instruments	Round One	Round One	Round One	Round Two	Round One
Exchange robotic instruments (including camera), with or without guided exchange	Round One	Round One	Round One	Round Two	Round One
Remove and clean camera when vision is obscured	Round One	Round One	Round One	Round One	Round One
Switch between left and right eye views while at the Patient Cart and Vision Cart	Round One	Round One	No Consensus	Round Two	No Consensus
Undock and insert trocars if they are pulled back too far	Round One	Round One	Round One	No Consensus	Round One
Insert new trocar or upsize trocar while the Patient Cart is docked	Round One	Round One	Round One	Round Two	Round One
Safely insert laparoscopic instruments	Round One	Round One	Round One	No Consensus	Round One
Manually release instruments <sup>a</sup>	Round Two	Round Two	Round Two	Round Two	No Consensus
Handle tissue with laparoscopic instruments without causing tissue injury or bleeding	Round One	Round One	Round One	No Consensus	Round One
Suction in the surgical field without causing tissue injury	Round One	Round One	Round One	No Consensus	Round One
Balance over- and under-suctioning	Round One	Round One	Round One	No Consensus	Round One
Perform controlled irrigation	Round One	Round One	Round One	No Consensus	Round One
Provide optimal, dynamic retraction	Round One	Round One	Round One	Round Two	Round One
Pass and remove sutures in the field	Round One	Round One	Round One	No Consensus	Round One
Cut sutures	Round One	Round One	Round One	No Consensus	Round One
Manipulate the uterus as needed during gynecological procedures	Round One	Round One	Round One	No Consensus	Round One
Manipulate the Foley catheter as needed during genitourinary procedures	Round One	Round One	Round One	No Consensus	No Consensus
Load and unload staplers <sup>a</sup>	Round Two	Round Two	No Consensus	Round Two	Round Two
Fire a manual stapler (linear or circular)	Round One	Round One	Round One	Round One	Round One
Provide direct pressure to control bleeding	Round One	Round One	Round One	No Consensus	Round Two
Remove specimens using a grasper	Round One	Round One	Round One	Round Two	Round One
Remove specimens by deploying a bag	Round One	Round One	Round One	Round Two	Round One
Undock the robot	Round One	Round One	Round One	Round One	No Consensus
Close fascial incisions	Round One	Round One	Round One	Round One	Round One
Close skin/subcutaneous incisions	Round One	Round One	Round One	Round Two	Round One

In round one and round two of the Delphi process, experts came to consensus on intra-operative technical skills that bedside assistants and other team members should have (green) or do not need (red) in a robotic operation. Experts did not come to consensus on the necessity of team members having a number of the skills (gray)

<sup>a</sup>Added in Round Two



**Table 3** Non-technical skill list (Color table online)

Skill	“Basic” bedside assistant (cases without assist port)	“Advanced” bedside assistant (cases with assist port)	Surgeon	Surgical technologist	Circulating nurse
State the surgical indication	No Consensus	Round One	Round One	Round Two	No Consensus
Describe the surgical steps	No Consensus	Round One	Round One	No Consensus	No Consensus
Confirm the patient’s medical appropriateness for the operation	No Consensus	No Consensus	Round One	Round One	No Consensus
State the common pitfalls of the operation and their management	No Consensus	Round Two	Round One	Round One	Round One
Identify commonly used robotic instruments by name and appearance	Round One	Round One	Round One	Round One	Round One
Troubleshoot CO2 insufflation when insufflation pressure is not as intended	Round One	Round One	Round One	Round One	Round One
Troubleshoot external collisions and adjust robotic arm positioning as needed	Round One	Round One	Round One	No Consensus	Round One
Troubleshoot if instrument does not load correctly	Round One	Round One	Round One	Round Two	Round One
Confirm appropriate instrument position (jaws are open and not clamped on tissue; wrist is straight) with surgeon prior to instrument exchange	Round One	Round One	Round One	Round Two	Round One
Describe the anatomy identified throughout the case	Round One	Round One	Round One	No Consensus	No Consensus
Troubleshoot robotic system faults	Round One	Round One	Round One	Round One	Round One
Describe the steps to convert to an open procedure	Round One	Round One	Round One	Round One	No Consensus
Identify instrument defects that may contribute to malfunction (e.g., unintentional cauterization due to damaged insulation)	Round One	Round One	Round One	Round One	Round Two
Perform closed-loop communication with the surgical team	Round One	Round One	Round One	Round One	Round One

In round one and round two of the Delphi process, experts came to consensus on non-technical skills that bedside assistants and other team members should have (green) or do not need (red) in a robotic operation. Experts did not come to consensus on the necessity of team members having a number of the skills (gray)

are prepared to participate in different cases. Such decisions can stem from skill rather than degree or prior time spent assisting.

There are multiple limitations within our study that moderate our findings. First, our developed list contains many items specific to the da Vinci surgical system. As emerging robotic surgical systems become available, additional work must generalize this list to reduce its platform specificity. Second, our experts included a limited number of surgical technologists and nurses as the study’s focus was on the skills required of bedside assistants. As such, experts may not have had the insight to appropriately determine the skills needed for each role. We defined expertise based on publications and experience conducting bedside assistant training, though other methods of expert selection may have led to different results. Additionally, 13 of our 14 experts practiced in the United States, which may limit the generalizability of these findings internationally. Similarly, the inclusion of an international expert, despite that expert’s prior experience publishing about bedside assistant training, may limit results given global practice pattern variation. Furthermore, there are also limitations to the application of our findings.

Experts did not come to consensus on a portion of the items. We opted to end the process after the second round as we perceived the lack of consensus at this point to reflect the quite varied practice patterns around the country and world. For example, experts seemed to identify quite disparate roles for surgical technologists. These different practice patterns may challenge the use of the list developed here. Future work could aim to further standardize role-specific expectations; nonetheless, focusing on standardizing the skills required of all bedside assistants may have broad applicability.

## Conclusion

We used the Delphi process to develop consensus guidelines around the skills required for bedside assistants in robotic surgery. These guidelines may serve as a framework for those designing training and assessments for bedside assistants. Standardized expectations and requirements for bedside assistant skills may allow for the consistency and quality needed in robotic surgery.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00464-024-11206-x>.

**Acknowledgements** We would like to acknowledge the 14 experts who participated in the study for their thoughtful and comprehensive review of all materials. We would also like to acknowledge Sherry Rogers, Melissa Boutte, Stacey Toplit, and Kristin Thompson for insight with early versions of the items and assistance connecting to appropriate resources.

**Funding** The authors received funding for this project through a Robotic Surgery Research Grant from the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and Intuitive Surgical.

## Declarations

**Disclosures** The authors received funding for this project through a Robotic Surgery Research Grant from the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and Intuitive Surgical. In addition, Riley Brian and Alyssa Murillo participate in the Intuitive-UCSF Simulation-Based Surgical Education Research Fellowship. Camilla Gomes participates in a research fellowship through Johnson & Johnson. Daniel Oh is part-time employed by Intuitive Surgical as a medical advisor. Hueylan Chern and Patricia O'Sullivan declare that they have no conflicts of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Britton CR, Francis I, Tay LJ, Krishnamoorthy B (2022) The role of the bedside assistant in robot-assisted surgery: a critical synthesis. *J Perioper Pract* 32(9):208–225
2. Onol FF, Sivaraman A, Andrich J. The Role of Bedside Assistant in Robotic Urological Surgery. In: Wiklund P, Mottrie A, Gundeti MS, Patel V, editors. *Robotic Urologic Surgery* [Internet]. Cham: Springer International Publishing; 2022 [cited 2024 Apr 13]. p. 39–49. Available from: [https://link.springer.com/https://doi.org/10.1007/978-3-031-00363-9\\_6](https://link.springer.com/https://doi.org/10.1007/978-3-031-00363-9_6)
3. Brian R, Oh D, Ifuku KA, Sarin A, O'Sullivan P, Chern H (2023) Experience matters for robotic assistance: an analysis of case data. *J Robot Surg* 17(5):2421–2426
4. Garbens A, Lay AH, Steinberg RL, Gahan JC (2021) Experienced bedside-assistants improve operative outcomes for surgeons early in their learning curve for robot assisted laparoscopic radical prostatectomy. *J Robot Surg* 15(4):619–626
5. Yu N, Saadat H, Finelli A, Lee JY, Singal RK, Grantcharov TP et al (2021) Quantifying the “Assistant Effect” in Robotic-Assisted Radical Prostatectomy (RARP): measures of technical performance. *J Surg Res* 260:307–314
6. Nayyar R, Yadav S, Singh P, Dogra P (2016) Impact of assistant surgeon on outcomes in robotic surgery. *Indian J Urol* 32(3):204
7. Lagrange F, Fiard G, Larose C, Eschwege P, Hubert J (2022) Role and training of the bedside surgeon in robotic surgery: a survey among french urologists-in-training. *Res Rep Urol* 14:17–22
8. Santos DA, Zhang L, Limmer AR, Gibson HM, Minetree C, Gollihar SH et al (2022) Protocolized training of advanced practice providers for robotic surgery improves the quality of intraoperative assistance. *JSLs* 26(3):e2022.00024
9. Moit H, Dwyer A, De Sutter M, Heinzel S, Crawford D (2019) A standardized robotic training curriculum in a general surgery program. *JSLs* 23(4):e2019.00045
10. Estes SJ, Goldenberg D, Winder JS, Juza RM, Lyn-Sue JR (2017) Best practices for robotic surgery programs. *JSLs* 21(2):e2016.00102
11. Schreuder H, Wolswijk R, Zweemer R, Schijven M, Verheijen R (2012) Training and learning robotic surgery, time for a more structured approach: a systematic review. *BJOG* 119(2):137–149
12. Stefanidis D, Huffman EM, Collins JW, Martino MA, Satava RM, Levy JS (2020) Expert consensus recommendations for robotic surgery credentialing. *Ann Surg* 276:88–93
13. Barnes KE, Brian R, Greenberg AL, Watanaskul S, Kim EK, O'Sullivan PS et al (2023) Beyond watching: harnessing laparoscopy to increase medical students' engagement with robotic procedures. *Am J Surg* S0002–9610(23):00092–00102
14. Greenberg AL, Syed SM, Alseidi A, O'Sullivan PS, Chern H (2022) Robotic training for medical students: feasibility of a pilot simulation curriculum. *J Robotic Surg* 17(3):1029–1038
15. Mullens CL, Van Horn AL, Marsh JW, Hogg ME, Thomay AA, Schmidt CR et al (2021) Development of a senior medical student robotic surgery training elective. *J Med Educ Curric Dev* 8:23821205211024070
16. Collins JM, Walsh DS, Hudson J, Henderson S, Thompson J, Zychowicz M (2021) Implementation of a standardized robotic assistant surgical training curriculum. *J Robot Surg* 16:789–797
17. Yuh B (2013) The bedside assistant in robotic surgery—keys to success. *Urol Nurs* 33(1):29–32
18. Association of Surgical Technologists. AST Guidelines for Best Practices on the Perioperative Role and Duties of the Surgical Technologist During Robotic Surgical Procedures [Internet]. 2017. Available from: [https://www.ast.org/uploadedFiles/Main\\_Site/Content/About\\_Us/ASTGuidelinesRoboticSurgicalProcedures.pdf](https://www.ast.org/uploadedFiles/Main_Site/Content/About_Us/ASTGuidelinesRoboticSurgicalProcedures.pdf)
19. Association of Surgical Assistants. Guideline Statement for the Surgical Assistant in Robotic Surgery [Internet]. 2017. Available from: [https://www.surgicalassistant.org/assets/docs/Guidelines\\_Surgical\\_Robotics.pdf](https://www.surgicalassistant.org/assets/docs/Guidelines_Surgical_Robotics.pdf)
20. Chen R, Rodrigues Armijo P, Krause C, Robotic Task Force SAGES, Siu KC, Oleynikov D (2020) A comprehensive review of robotic surgery curriculum and training for residents, fellows, and postgraduate surgical education. *Surg Endosc* 34(1):361–7
21. Nasa P, Jain R, Juneja D (2021) Delphi methodology in healthcare research: how to decide its appropriateness. *World J Methodol* 11(4):116–129
22. Humphrey-Murto S, Varpio L, Wood TJ, Gonsalves C, Uffholz LA, Mascioli K et al (2017) The use of the delphi and other consensus group methods in medical education research: a review. *Acad Med* 92(10):1491–1498
23. Brian R, Murillo A, Oh D, Chern H, O'Sullivan P (2024) Comparing observed and preferred instruction in robotic surgery. Under Rev. <https://doi.org/10.1016/j.surg.2024.06.043>
24. Schiff L, Tsafir Z, Aoun J, Taylor A, Theoharis E, Eisenstein D (2016) Quality of communication in robotic surgery and surgical outcomes. *JSLs* 20(3):201600026

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.