



解放军医学杂志

Medical Journal of Chinese People's Liberation Army

ISSN 0577-7402,CN 11-1056/R

《解放军医学杂志》网络首发论文

题目： 脓毒症性凝血病诊疗中国专家共识
作者： 宋景春，丁仁戡，吕奔，梅恒，王岗，张伟，周静，郭军
收稿日期： 2024-08-02
网络首发日期： 2024-09-24
引用格式： 宋景春，丁仁戡，吕奔，梅恒，王岗，张伟，周静，郭军. 脓毒症性凝血病诊疗中国专家共识[J/OL]. 解放军医学杂志.
<https://link.cnki.net/urlid/11.1056.R.20240923.1715.002>



网络首发：在编辑部工作流程中，稿件从录用到出版要经历录用定稿、排版定稿、整期汇编定稿等阶段。录用定稿指内容已经确定，且通过同行评议、主编终审同意刊用的稿件。排版定稿指录用定稿按照期刊特定版式（包括网络呈现版式）排版后的稿件，可暂不确定出版年、卷、期和页码。整期汇编定稿指出版年、卷、期、页码均已确定的印刷或数字出版的整期汇编稿件。录用定稿网络首发稿件内容必须符合《出版管理条例》和《期刊出版管理规定》的有关规定；学术研究成果具有创新性、科学性和先进性，符合编辑部对刊文的录用要求，不存在学术不端行为及其他侵权行为；稿件内容应基本符合国家有关书刊编辑、出版的技术标准，正确使用和统一规范语言文字、符号、数字、外文字母、法定计量单位及地图标注等。为确保录用定稿网络首发的严肃性，录用定稿一经发布，不得修改论文题目、作者、机构名称和学术内容，只可基于编辑规范进行少量文字的修改。

出版确认：纸质期刊编辑部通过与《中国学术期刊（光盘版）》电子杂志社有限公司签约，在《中国学术期刊（网络版）》出版传播平台上创办与纸质期刊内容一致的网络版，以单篇或整期出版形式，在印刷出版之前刊发论文的录用定稿、排版定稿、整期汇编定稿。因为《中国学术期刊（网络版）》是国家新闻出版广电总局批准的网络连续型出版物（ISSN 2096-4188，CN 11-6037/Z），所以签约期刊的网络版上网络首发论文视为正式出版。

指南与共识

[作者简介] 宋景春，医学博士，主任医师，主要从事血栓与止血危重病方面的研究

[通信作者] 宋景春，E-mail: songjingchun@126.com

脓毒症性凝血病诊疗中国专家共识

宋景春^{1*}，丁仁彧²，吕奔³，梅恒⁴，王岗⁵，张伟⁶，周静⁷，郭军⁸，中国医药教育协会血栓与止血危重病专业委员会，全军重症医学专业委员会

¹解放军联勤保障部队第908医院重症医学科，江西南昌 330002；²中国医科大学附属第一医院重症医学科，辽宁沈阳 110001；³中南大学湘雅二医院血液科，湖南长沙 410011；⁴华中科技大学同济医学院附属协和医院血液科，湖北武汉 430030；⁵西安交通大学第二附属医院重症医学科，陕西西安 710004；⁶解放军联勤保障部队第900医院急诊科，福建福州 350002；⁷四川大学华西医院实验医学科，四川成都 610041；⁸四川大学华西医院重症医学科，四川成都 610041

[摘要] 脓毒症性凝血病(SIC)是脓毒症导致的内皮损伤和凝血紊乱，是导致脓症患者死亡的重要因素。对SIC实现早期诊断和治疗有望改善脓毒症患者的预后。2019年国际血栓止血学会(ISTH)颁布了首部SIC诊疗指南，但我国尚无相关规范。因此，中国医药教育协会血栓与止血危重病专业委员会和全军重症医学专业委员会共同编写了《脓毒症性凝血病诊疗中国专家共识》。该共识包括发病机制、分型、实验室评估、诊断和治疗等5个部分，共14条推荐意见，以便指导相应的临床工作。

[关键词] 脓毒症；凝血病；免疫血栓；血栓炎症；表型；抗凝

[中图分类号] R459.7 [文献标志码] A [DOI] 10.11855/j.issn.0577-7402.1189.2024.0918

[收稿日期] 2024-08-02

[录用日期] 2024-09-08

Chinese expert consensus on diagnosis and treatment of sepsis-induced coagulopathy

Song Jing-Chun^{1*}, Ding Ren-Yu², Lü Ben³, Mei Heng⁴, Wang Gang⁵, Zhang Wei⁶, Zhou Jing⁷, Guo Jun⁸; Chinese Society of Thrombosis, Hemostasis and Critical Care, Chinese Medicine Education Association; People's Liberation Army Professional Committee of Critical Care Medicine

¹Department of Critical Care Medicine, the 908th Hospital of the PLA Joint Logistic Support Force, Nanchang, Jiangxi 330002, China

²Department of Critical Care Medicine, the First Affiliated Hospital of Chinese Medical University, Shenyang, Liaoning 110001, China

³Department of Hematology, Xiangya Second Hospital, Central South University, Changsha, Hunan 410011, China

⁴Department of Hematology, Union Hospital Affiliated to Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430030, China

⁵Department of Critical Care Medicine, the Second Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaanxi 710004, China

⁶Department of Emergency, the 900th Hospital of the PLA Joint Logistic Support Force, Fuzhou, Fujian 350002,

China

⁷Department of Laboratory Medicine, ⁸Department of Critical Care Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan 610041, China

*Corresponding author; E-mail: songjingchun@126.com

[Abstract] Sepsis-induced coagulopathy (SIC), a critical condition stemming from sepsis that leads to endothelial injury and coagulation dysregulation, is a pivotal determinant in the mortality of sepsis patients. The timely diagnosis and therapeutic intervention for SIC are anticipated to enhance the clinical outcomes for these patients. In 2019, the International Society on Thrombosis and Haemostasis (ISTH) introduced the inaugural guidelines for managing SIC, yet China has not established equivalent protocols. In response, Chinese Society of Thrombosis, Hemostasis and Critical Care, Chinese Medicine Education Association, in collaboration with Chinese People's Liberation Army Professional Committee of Critical Care Medicine, has developed this consensus document. It is comprehensively structured into five distinct sections: etiology, classification, evaluative approaches, diagnosis and therapeutic strategies, encompassing a total of 14 evidence-based recommendations aimed at guiding clinical practice.

[Keywords] sepsis; coagulopathy; immunothrombosis; thromboinflammation; phenotypes; anticoagulation

脓毒症是机体对感染的反应失调导致的致死性器官功能障碍^[1]。脓毒症性凝血病(sepsis-induced coagulopathy, SIC)是脓毒症导致的血管内皮细胞损伤和凝血紊乱^[2]。据统计,全世界有 24.0%~60.0%的脓毒症患者发生 SIC, 而我国则高达 67.9%, 如处理不当可发展为弥散性血管内凝血(disseminated intravascular coagulation, DIC), 使患者病死率增高 2 倍^[3-7]。国际血栓止血学会(International Society on Thrombosis and Haemostasis, ISTH)和欧洲心脏病学会(European Society of Cardiology, ESC)已发布了 SIC 的诊断标准和治疗指南, 但迄今为止我国尚未制定 SIC 的诊疗规范^[8-9]。因此, 中国医药教育协会血栓与止血危重病专业委员会和全军重症医学专业委员会共同编写了本共识, 内容包括发病机制、分型、实验室评估、诊断和治疗等 5 个部分, 共 14 条推荐意见, 推荐强度及循证证据等级见表 1、2, 以供临床医护人员参考。

表 1 临床推荐强度分级

Tab. 1 Classification and description of clinical recommendation

推荐强度	等级释义及临床建议
I	强。循证证据肯定或良好(A-B 级); 循证证据一般(C-D 级), 但在国内外指南中明确推荐, 能够改善健康结局, 利大于弊
II	中等。循证证据一般(C-D 级); 可改善健康结局
III	弱。循证证据不足或矛盾; 无法明确利弊, 但可能改善健康结局

表 2 循证证据等级


Tab. 2 Evidence-based level

证据等级	分级释义
A	基于多个随机对照试验的荟萃分析或系统评价; 大样本随机对照试验
B	基于至少一个质量较高的随机对照试验; 设计规范、结果明确的观察性研究或横断面研究; 前瞻性队列研究
C	基于设计良好的非随机病例对照研究, 观察性研究, 非前瞻性队列研究
D	基于非随机的回顾性研究; 病例报告; 专家共识

1 发病机制

推荐意见 1 SIC 是从适度免疫血栓发展成过度血栓炎症的过程(推荐强度 II, 证据等级 D)

免疫血栓是感染早期机体在微血管内形成的具有免疫功能的血栓。免疫血栓过度形成可导致血栓炎症的失调和扩散, 进而发生 SIC^[10]。免疫血栓形成是由白细胞和血小板共同参与的先天固有免疫反应。中性粒细胞、单核细胞和巨噬细胞等表面均可表达各种受体, 如 Toll 样受体、Fc γ 受体、G 蛋白偶联受体、黏附受体和细胞因子受体, 以发现病原体的入侵。中性粒细胞死亡后释放出 DNA 和组蛋白组成中性粒细胞胞外诱捕网(neutrophil extracellular traps, NETs), NETs 与血小板、纤维蛋白原共同组成免疫血栓, 对病原体进行拦截、分隔、捕捉和灭杀。存活的中性粒细胞还能以非细胞溶解的方式排出组蛋白和线粒体 DNA, 参与 NETs 的形成^[11]。NETs 可为血管性血友病因子(von Willebrand factor, vWF)、纤维蛋白原、凝血因子XII、组织因子(tissue factor, TF)等促凝因子和携带 TF 等促凝因子的细胞外囊泡(extracellular vesicles, EV)提供支架, 以促进血栓形成^[12]。巨噬细胞和单核细胞也可通过类似中性粒细胞的机制形成胞外诱捕网(extracellular traps, ETs), 加强免疫血栓形成效应^[13]。当系统性免疫反应持续进行时, 广泛形成的免疫微血栓会导致血栓炎症失控, 加重血管内皮细胞功能障碍, 形成毛细血管渗漏和消耗性凝血病, 乃至 DIC 和多器官功能衰竭^[14](图 1)。



注: 黑色线条代表免疫血栓形成过程, 蓝色线条代表血栓炎症形成过程

图 1 免疫血栓与血栓炎症的发生机制


Fig.1 Mechanism of immunothrombosis and thromboinflammation development

2 分型

推荐意见 2 SIC 可分为高凝血症和低凝血症两种表型, 临床上可表现为血栓形成和(或)出血(推荐强度 II, 证据等级 C)

SIC 既可表现为以血栓形成为主的高凝血症, 又可呈现为以出血倾向为主的低凝血症^[15-16](图 2)。当病原体作用于机体后, 可通过病原相关分子模式(pathogen-associated molecular patterns, PAMPs)或损伤相关分子模式(damage-associated molecular patterns, DAMPs)造成内皮细胞损伤, 协同炎症细胞和血小板共同释放大量 TF, 通过外源性凝血途径激活凝血酶, 促进纤维蛋白与血小板在微血管内广泛形成微血栓, 最终导致多器官功能损害^[17]。脓毒症时中性粒细胞和单核细胞可通过含半胱氨酸的天冬氨酸蛋白水解酶 11(cysteiny aspartate specific proteinase-11, caspase-11)介导的细胞焦亡途径和核苷酸结合寡聚化结构域样受体热

蛋白结构域相关蛋白 3(nucleotide oligomerization domain-like receptor thermal protein domain associated protein 3, NLRP3)炎性小体促进 TF 大量活化,这一过程也被称为 TF“解密”^[18-19]。内皮细胞损伤还会抑制抗凝血酶(antithrombin, AT)、活化蛋白 C(activated protein C, APC)系统和组织因子途径抑制剂(tissue factor pathway inhibitor, TFPI)等生理性抗凝物质的合成与释放,增强纤溶酶原激活物抑制剂-1(plasminogen activator inhibitor-1, PAI-1)、凝血酶激活的纤溶抑制物(thrombin-activatable fibrinolysis inhibitor, TAFI)活性,进而促进血栓的形成。当凝血因子和血小板被大量消耗后,可导致显著的低凝血症,机体表现为出血倾向甚至发生出血^[20]。



注: SIC. 脓毒症性凝血病

图 2 SIC 的表型分类

Fig.2 Phenotype classification of sepsis-induced coagulopathy

3 实验室评估

推荐意见 3 充分使用凝血实验室指标对 SIC 患者的凝血功能进行动态评估(推荐强度 II, 证据等级 C)

根据检测原理的不同,凝血实验室指标可分为以血浆标本为检测对象的指标和以全血标本为检测对象的指标。血浆标本的凝血指标重在反映某一个或几个凝血蛋白的活性,全血标本的凝血指标重在反映凝血功能的整体效应。基于 SIC 的病理生理机制,凝血实验室指标改变可呈现典型的时相特征(表 3)。

表 3 SIC 不同时相的凝血实验室指标改变

Tab.3 Changes in coagulation laboratory indicators at different phases of SIC

项目	普通脓毒症	SIC 高凝血症	SIC 低凝血症	DIC
血小板计数	升高/减少	升高/减少	减少	减少
PT	正常	正常或缩短	延长	延长
APTT	正常	正常或缩短	延长	延长
纤维蛋白原	正常	正常或升高	降低	降低
D-二聚体	升高	升高	升高	升高
纤维蛋白降解产物	升高	升高	升高	升高
内皮细胞功能	正常或升高	升高	升高	升高
t-PAIC	正常	正常	正常或升高	升高
凝血酶	正常或升高	升高	升高	升高
纤溶酶	正常	正常	正常或升高	升高
血栓弹力图	R 时间	正常	缩短	延长

	K 时间	正常	缩短	延长	延长
	α 角	正常	增大	减小	减小
	血块最大振幅	正常	增大	减小	减小
	活化凝血时间	正常	缩短	延长	延长
凝血与血小板功能分析	凝血速率	正常	升高	降低	降低
	血小板功能	正常	升高	降低	降低

注: PT. 凝血酶原时间; APTT. 活化部分凝血活酶时间; t-PAIC. 纤溶酶原激活物-纤溶酶原激活抑制物-1 复合物; SIC. 脓毒症性凝血病; DIC. 弥散性血管内凝血

SIC 高凝血症时, 血管内皮细胞出现损伤, 血浆血栓调节蛋白(thrombomodulin, TM)水平升高; 内皮细胞损伤后, 凝血酶活化, 凝血酶-抗凝血酶复合物(thrombin-antithrombin complex, TAT)水平可出现升高, 凝血酶原时间(prothrombin time, PT)和活化部分凝血活酶时间(activated partial thromboplastin time, APTT)可处于正常范围或轻度缩短; 微血栓形成后可继发纤溶活动, 进而出现 D-二聚体和纤维蛋白降解产物(fibrin degradation products, FDP)水平升高; 在炎症反应的刺激下, 部分患者的血小板计数可出现增多甚至轻度减少; 作为急性反应时相蛋白, 纤维蛋白原水平也可升高^[21]。血栓弹力图(thromboelastograph, TEG)可出现 R 时间(代表凝血因子活性)缩短, K 时间(代表纤维蛋白原功能)缩短, α 角(代表纤维蛋白原功能)增大, 血块最大振幅(maximum amplitude, MA, 代表血小板功能)增大, 凝血指数(coagulation index, CI)增高, 整体呈现高凝状态^[22]。凝血与血小板功能分析可发现活化凝血时间(activated clotting time, ACT, 代表凝血因子活性)缩短, 凝血速率(clotting rate, CR, 代表纤维蛋白原功能)增高, 以及血小板功能(platelet function, PF)增强^[23]。此时患者可无特征性的临床表现, 但可出现器官功能损害或凝血功能的实验室指标异常。如患者存在糖尿病、冠心病等易导致血栓形成的基础疾病, 则脓毒症可能诱发心肌梗死、脑梗死等缺血事件^[24]。

SIC 低凝血症时, 血管内皮细胞损伤加重, 凝血酶过度活化, 血浆 TM 和 TAT 水平均明显升高; 凝血底物过度消耗, 可导致血小板计数减少, PT、APTT 明显延长, 纤维蛋白原水平下降; 促进纤溶活动的组织型纤溶酶原激活物(tissue plasminogen activator, t-PA)和 PAI-1 水平均升高, 但 PAI-1 升高的幅度更大, 血浆 t-PA/PAI-1 比值降低, 总体呈现纤溶抑制状态。此时 D-二聚体和 FDP 水平会明显升高, 但纤溶酶-抗纤溶酶复合物(plasmin-antiplasmin complex, PIC)水平升高不明显。出现脓毒症性休克时, 组织缺血缺氧造成血管内皮细胞损伤进一步加重, 纤溶酶原激活物-纤溶酶原激活抑制物-1 复合物(tissue plasminogen activator-plasminogen activator inhibitor-1 complex, t-PAIC)水平可明显升高^[25]。TEG 可出现 R 时间和 K 时间延长, α 角减小, MA 和 CI 降低, 整体呈现低凝状态^[26-27]。凝血与血小板功能分析可发现 ACT 延长、CR 降低和 PF 减弱。此时, 脓毒症患者可因广泛血栓形成出现多器官功能衰竭、肢端发绀甚至坏疽, 如凝血底物严重消耗可合并皮肤淤斑、牙龈出血、消化道出血、尿道出血等出血表现, 同时 PIC 水平明显升高。

4 诊断

推荐意见 4 推荐使用中国 SIC 评分系统诊断 SIC, 使用 ISTH-DIC 评分诊断脓毒症性 DIC(推荐强度 II, 证据等级 B)

DIC 是因致病因素诱发机体产生促凝物质激活凝血酶, 导致广泛形成微血栓, 凝血因子大量消耗并继发纤溶亢进, 引起全身出血与微循环衰竭的综合征^[28]。2001 年, ISTH 将 DIC 分为显性 DIC(即失代偿期 DIC)与非显性 DIC(即代偿期 DIC), 强调在 DIC 代偿期需每日评估 DIC 是否进展到失代偿期, 并推荐采用 DIC 诊断积分系统诊断显性 DIC^[29]。近年研究发现, 在 DIC 代偿期即开始针对 DIC 进行治疗可能使患者获益^[30]。因此, ISTH 科学标准化委员会于 2019 年提出使用 SIC 诊断脓毒症性 DIC 的代偿期^[31]。

ISTH 推荐的 SIC 评分系统纳入了 PT 的国际标准化比值(international normalized ratio,

INR)、血小板计数和脓毒症相关性器官功能衰竭评价(sepsis-related organ failure assessment, SOFA)3项指标,评分 ≥ 4 分即可诊断 SIC(表 4)^[31]。ISTH 推荐的 DIC 积分系统纳入了 PT、血小板计数、纤维蛋白原和 D-二聚体等指标进行积分,评分 ≥ 5 分即可诊断显性 DIC^[29]。当 ISTH-DIC 积分达到最高的 8 分且血乳酸水平 >2 mmol/L 时,可诊断凝血衰竭^[32]。已有研究显示, SIC 诊断标准对死亡的预测敏感度要高于 ISTH-DIC 诊断标准^[33]。遗憾的是, ISTH-SIC 诊断方案中血小板计数是以 $<150 \times 10^9/L$ 开始积分的,符合欧美人群血小板计数以 $<150 \times 10^9/L$ 作为血小板减少的诊断标准,而中国人群以血小板计数 $<100 \times 10^9/L$ 作为血小板减少的诊断标准。因此,直接使用 ISTH-SIC 诊断标准会导致中国人群诊断 SIC 的假阳性率增高,并导致部分血小板计数在 $(100 \sim 150) \times 10^9/L$ 范围内的 SIC 患者接受不必要的抗凝治疗^[34]。依据《凝血障碍诊断规范》,本共识推荐应用中国 SIC 诊断标准诊断 SIC,使用 ISTH-DIC 评分诊断脓毒症性 DIC^[32]。


表 4 SIC 与 ISTH-DIC 诊断标准

Tab.4 Diagnostic criteria for SIC and ISTH-DIC

指标	ISTH-SIC	中国 SIC	ISTH-DIC	凝血衰竭	评分
SOFA(分)	1	1	-	-	1
	≥ 2	≥ 2	-	-	2
血小板($\times 10^9/L$)	≤ 150	≤ 100	≤ 100	-	1
	≤ 100	≤ 50	≤ 50	≤ 50	2
PT 延长值(s)/INR	(1.2, 1.4]	(1.2, 1.4]	[3, 6)	-	1
	>1.4	>1.4	≥ 6	≥ 6	2
D-二聚体($\mu g/mL$)	-	-	[2.5, 5.0)	-	2
	-	-	≥ 5.0	≥ 5.0	3
纤维蛋白原(g/L)	-	-	< 1.0	< 1.0	1
乳酸(mmol/L)	-	-	-	>2	
总分(分)	≥ 4	≥ 4	≥ 5	-	

注: ISTH 国际血栓止血学会; SIC. 脓毒症性凝血病; DIC. 弥散性血管内凝血; SOFA. 脓毒症相关性器官功能衰竭评价; PT. 凝血酶原时间; INR. 国际标准化比值

该诊断方案为 SIC 的早期诊断和抗凝治疗时机的选择提供了依据,有利于临床实施。但仍存在以下局限:(1)SIC 诊断标准主要用于诊断处于脓毒症性 DIC 代偿期的患者,并不能区分 SIC 患者的血液高凝或低凝状态;(2)ISTH-DIC 患者积分最高值为 8 分,不能有效区分 DIC 患者凝血障碍的严重程度。因此,本共识依据《凝血障碍诊断规范》,推荐应用中国 SIC 诊断标准诊断 SIC,使用 ISTH-DIC 评分诊断脓毒症性 DIC,使用凝血衰竭诊断标准诊断特重型 DIC(图 3)^[27]。




注: SIC. 脓毒症性凝血病; DIC. 弥散性血管内凝血

图 3 SIC 诊断关系图

Fig.3 Diagnostic diagram of SIC

5 治疗

SIC 的治疗主要遵循“三抗一防”的原则，即抗凝、抗炎、抗感染和预防稀释性凝血病(图 4)。



注：DIC. 弥散性血管内凝血；SIC. 脓毒症性凝血病

图 4 SIC 的“三抗一防”治疗时机

Fig.4 Timing of 'three antis and one prevention' treatment for SIC


推荐意见 5 积极有效的抗感染是治疗 SIC 的前提(推荐强度 I，证据等级 B)

感染是导致 SIC 的根本原因，因此积极有效的抗感染是成功治疗 SIC 的前提。具体步骤包括及时全面筛查感染源、控制感染源和尽早合理使用抗生素^[35-37]。控制感染源的具体措施包括脓肿引流、感染坏死组织清创、移除潜在感染装置或对持续性微生物污染源进行确定性控制等^[38]。在进行相关有创操作前，还要对 SIC 患者的出血风险进行评估^[39]。

此外，尽早合理使用抗生素进行抗感染治疗时，需关注部分抗生素对凝血功能的影响。已有文献报道，头孢哌酮、利福平和哌拉西林他唑巴坦等广谱抗生素可破坏肠道菌群并抑制维生素 K 的合成，导致维生素 K 相关的凝血因子缺乏，凝血实验室检测表现为 PT 和 APTT 延长^[40-42]。一旦因使用抗生素导致维生素 K 相关的凝血因子缺乏，除停用相关抗生素外，如无出血可补充维生素 K，如合并出血还需使用凝血酶原复合物。替加环素相关的凝血障碍可表现为 PT 和 APTT 延长、纤维蛋白原水平降低、血小板减少；其机制除了抑制维生素 K 合成外，还包括影响肝功能和凝血因子合成，因此停用替加环素后仅补充维生素 K 难以改善凝血指标，通常还需输注血制品^[43]。部分抗生素能够抑制骨髓合成血小板而导致血小板减少，如利奈唑胺，相应的治疗为停用抗生素或联合使用促血小板生成药物^[44]。此外，抗生素引起的免疫性血小板减少往往进展迅速，严重者血小板计数可 $<10 \times 10^9/L$ ，甚至造成严重出血。已有报道可引起免疫性血小板减少的抗生素包括氨曲南、万古霉素、哌拉西林、达托霉素、甲氧苄啶-磺胺甲恶唑和利福平^[45]。治疗措施除停用相关抗生素外，还需采用激素或丙种球蛋白冲击治疗，必要时需行血浆置换^[46]。

推荐意见 6 推荐对 SIC 患者进行抗炎治疗(推荐强度 II，证据等级 C)

脓毒症从本质上来说是一种由宿主免疫反应介导的炎症性疾病，SIC 的发生发展与系统性炎症失调密切相关。因此，抗炎治疗有望治疗脓毒症并减轻凝血紊乱的严重程度。糖皮质激素、免疫球蛋白、维生素 C、乌司他丁和血液净化是目前研究较多的抗炎疗法，但均缺乏充分有效的循证医学依据^[47](图 5)。



注：SIC. 脓毒症性凝血病

图5 SIC 的抗炎治疗方法

Fig.5 Anti-inflammatory Treatment for SIC

糖皮质激素是目前临床上应用最广泛的抗炎药物。1960年，Bennett等^[48]进行了第一个尝试使用糖皮质激素治疗脓毒症的临床试验。遗憾的是，这项试验并未显示糖皮质激素有效。2018年，Rochweg等^[49]对包括10 194例患者的42项随机对照试验进行荟萃分析，结论显示低剂量的糖皮质激素可能有益。同年，Venkatesh等^[50]对3800例患者进行的随机对照试验结果显示，接受机械通气的脓毒症休克患者行氢化可的松持续输注治疗，与安慰剂相比并不影响其90 d病死率。因此，2021版国际脓毒症指南仅推荐脓毒症休克且需血管加压素维持血压的成年脓症患者使用低剂量氢化可的松^[51]。2024年美国 and 欧洲重症学会共同发布的应用糖皮质激素治疗脓毒症的指南也指出，小剂量短期使用糖皮质激素可小幅度降低病死率，并逆转器官功能障碍^[52]。

免疫球蛋白也是临床常用的免疫调节药物，具有抑制内皮细胞表达黏附分子和趋化因子、抑制补体过度活化、中和毒素、减轻毛细血管渗漏和保护血管内皮的作用^[53-54]。2015年，Ishikura等^[55]报道了一项应用丙种球蛋白治疗41例脓毒症合并凝血障碍患者的回顾性研究，结果显示接受丙种球蛋白治疗(5.0 g/d，共3 d)的脓症患者凝血指标(PT-INR下降、APTT缩短、PAI-1水平降低)、炎症指标(C反应蛋白和降钙素原水平降低)均明显改善，病死率也明显降低(5.3% vs. 18.2%)。2016年，Wand等^[56]报道应用IgM免疫球蛋白治疗严重脓症患者可改善血小板减少与低纤维蛋白原血症。后续的荟萃分析结果显示，IgG和IgM(RR=0.69, 95% CI 0.55-0.85)可降低患者的病死率，但因证据质量较低，并未得到2021版国际脓毒症指南的明确推荐^[51,57-58]。需要注意的是，部分品牌的药物说明书提示应用免疫球蛋白可能导致血栓栓塞，其机制与药品生产过程中混有凝血因子有关^[59]。

维生素C又称抗坏血酸，是一种水溶性维生素。因为人体无法合成维生素C，所以其获取主要依靠饮食摄入。维生素C具有抗氧化、改善毛细血管通透性、减轻毛细血管损伤和抗菌等的药理学作用^[60-61]。已有研究显示，危重患者的血浆维生素C水平明显下降，故有学者认为补充外源性维生素C有望治疗脓毒症^[62-63]。2017年，一项单中心回顾性研究报告，在对脓毒症和脓毒症休克患者联合使用高剂量维生素C、氢化可的松和硫酸胺素后，血管加压素治疗时间缩短，病死率明显降低(8.5% vs. 40.4%)^[64]。但多项RCT研究及荟萃分析结果均显示，使用维生素C、氢化可的松和硫酸胺素的组合治疗对病死率并无明显改善^[65-68]。因此，2021版国际脓毒症指南暂未推荐通过静脉注射维生素C治疗脓毒症^[51]。


乌司他丁是一种主要由肝细胞产生、存在于人类尿液或血液中的胰蛋白酶抑制剂^[69]。乌司他丁可通过降低p38丝裂原活化蛋白激酶(p38-MAPK)的磷酸化和核因子-κB(NF-κB)活化发挥抗炎作用，并能抑制中性粒细胞释放弹性蛋白酶和炎性介质^[70-71]。一项纳入13项单中心随机对照试验的荟萃分析显示，乌司他丁可降低脓症患者血清白细胞介素-6(IL-6)水平，

并明显降低脓毒症患者的全因死亡率，但迄今尚无多中心随机对照研究的相关证据^[72]。

肾脏替代治疗(renal replacement therapy, RRT)已被广泛应用于脓毒症合并急性肾损伤患者的治疗，而近年来研究认为其还具有抗炎治疗的作用^[73]。具体机制包括：(1)清除细胞因子：已有研究证实，高水平的细胞因子与脓毒症相关器官损伤严重程度和病死率增高有关^[74]。标准 RRT 治疗能够有效清除分子量在 40 kD 以下的细胞因子，包括血浆 IL-1 β 、IL-1 α 、IL-2、IL-4、IL-6、IL-8、IL-10、 γ 干扰素(interferon- γ , IFN- γ)和肿瘤坏死因子等^[75-76]。(2)减轻组织水肿：因为 RRT 治疗可实现精准的容量管理，从而减轻组织水肿，改善血管通透性，促进淋巴回流。(3)稳定内环境：血液净化能够纠正电解质紊乱和酸中毒，维持内环境稳定，改善凝血因子活性^[77]。(4)减少凝血酶活化：RRT 可通过降低细胞因子水平，减轻内皮细胞损伤，减少凝血酶活化，从而实现间接的抗凝作用^[78]。一项探讨连续性 RRT(continuous RRT, CRRT)对脓症患者凝血功能影响的单中心随机对照研究显示，经 CRRT 治疗 7 d 后常规凝血指标 PT、APTT、血浆纤维蛋白原和 D-二聚体水平均较常规治疗组更接近正常范围，炎症指标 C 反应蛋白、肿瘤坏死因子和 IFN- γ 水平降幅也更大^[79]。另一项小样本量研究对比分析了 CRRT 与间断血液透析治疗对脓毒症合并肾损伤患者凝血功能和炎症因子水平的影响，结果显示，与间断血液透析相比，接受 CRRT 治疗的患者 IL-6、IL-8、超敏 C 反应蛋白、肿瘤坏死因子 α 、PT、APTT、纤维蛋白原和血小板计数均优于普通血液透析组($P<0.05$)^[80]。此外，采用多黏菌素 B 血液吸附治疗能够去除脓症患者体内的内毒素，也被认为是脓毒症的有效治疗方法^[81]。JSEPTIC-DIC^[82]和 EUPHRATES^[83]试验结果显示，多黏菌素 B 血液吸附可改善血浆内毒素水平明显升高、PT-INR >1.4 或乳酸 >3 mmol/L 的脓毒症患者的预后，但对所有脓毒症患者的全因死亡率无明显影响^[84]。

推荐意见 7 当患者达到 SIC 诊断标准时应尽早启动抗凝治疗(推荐强度 II, 证据等级 C)

准确把握 SIC 患者的抗凝时机是精准施治的基本要求。根据抗凝治疗目的的不同，SIC 患者的抗凝指征有两种情况。一是当脓症患者达到 SIC 诊断标准时启动抗凝^[31](图 6)。此时启动抗凝的目的是保护血管内皮、减少凝血底物消耗、抑制血栓形成并改善器官血供。二是脓症患者出现明确的血栓并发症时需启动抗凝^[51]。此时启动抗凝的目的是抑制血栓蔓延，以利于血栓自溶和管腔再通。需要注意的是，活动性出血是抗凝治疗的禁忌证。此外，治疗脓毒症时并非一发现高凝状态就要抗凝，因为脓毒症早期形成的免疫血栓具有杀灭病原体和防止病原体扩散的作用^[85]，过早启动抗凝可能导致感染加重。一项针对日本 42 家 ICU 的 2663 例脓毒症患者的回顾性分析发现，抗凝治疗仅能改善 SIC 患者的预后，而无法使所有脓症患者获益^[86]。



注：SIC. 脓毒症性凝血病；DIC. 弥散性血管内凝血

图 6 SIC 的抗凝治疗药物选择

Fig.6 Selection of anticoagulant drugs for SIC treatment

SIC 患者往往合并器官功能障碍，易对抗凝药物的代谢造成影响，故应选择胃肠外抗凝

药物。国内目前常用的抗凝药物包括肝素类药物、直接凝血酶抑制剂和丝氨酸蛋白酶抑制剂(表 5)。SIC 患者的抗凝药物选择在日本作为一线抗凝药物的重组人 TM 和抗凝血酶(recombinant human antithrombin, rhAT), 在我国尚未供应, 故本文不做推荐^[87]。此外, 因为 SIC 患者本身存在凝血障碍, 所以选择抗凝药物前需先评估其出血风险(图 6)。



表5 SIC 抗凝药物特点

Tab.5 Characteristics of anticoagulant for SIC patients

药物分类	药物名称	半衰期	代谢途径	初始剂量	给药方式	监测方法	并发症	推荐意见
肝素类药物	普通肝素	60~90 min	肾脏	1~4 U/(kg·h)	静脉注射	(1)TEG 肝素酶对比试验: R 普通检测/R 肝素酶检测的比值为 1.5~2.0; (2)抗 Xa 活性 0.5~0.7 U/mL; (3)APTT 延长至基线值的 1.5 倍以内	出血、肝素诱导的血小板减少症	推荐
	低分子肝素	2.0~4.5 h	肾脏	50 U/kg, 每 12 h 一次	皮下或静脉注射	(1)皮下注射后 4 h 或静脉注射后 10 min 的抗 Xa 活性为 0.3~0.5 U/mL; (2)动态监测抗 Xa 活性或 TEG-R 时间	出血、肝素诱导的血小板减少症	推荐
	磺达肝癸钠	17~21 h; 老年人和肾功能不全者延长	肾脏	1.5~2.5 mg, 1 次/d	皮下注射	抗 Xa 活性或 TEG-R 时间	出血、恶心、呕吐、肝酶升高	不推荐
丝氨酸蛋白酶抑制剂	甲磺酸奈莫司他	5~8 min	80%经肝脏, 20%由红细胞降解	2~20 mg/h	静脉注射	APTT 或 TEG-R 时间	高钾血症、出血	推荐
直接凝血酶抑制剂	阿加曲班	39~51 min	肝脏代谢	1~2 μg/(kg·min)	静脉注射	APTT 或 TEG-R 时间至基线值的 1.5 倍以内	出血、肝功能异常	不推荐
	比伐卢定	25 min	80%蛋白水解, 20%经肾脏	1~8 μg/(kg·min)	静脉注射	APTT 或 TEG-R 时间至基线值的 1.5 倍以内	疼痛、低血压、恶心	不推荐

注: SIC. 脓毒性凝血病; TEG. 血栓弹力图; APTT. 活化部分凝血活酶时间

推荐意见 8 SIC 患者抗凝药物宜首选肝素类药物(推荐强度II, 证据等级 C)

普通肝素的半衰期短、监测方便,一旦过量可用鱼精蛋白中和,故常作为 SIC 患者抗凝的首选药物^[88]。普通肝素常选择静脉给药,决定药物剂量时需兼顾出血风险与抗凝效果。SIC 患者使用肝素抗凝可从小剂量[1~4 U/(kg·h)]开始,根据实验室指标进行滴定式治疗。APTT 是最常用于监测普通肝素抗凝效果的指标,通常以 APTT 延长至基线值的 1.5 倍为宜。但因为 SIC 患者本身可能存在 APTT 延长,再用 APTT 指导普通肝素抗凝会增加出血风险,故可使用血栓弹力图肝素酶对比试验来指导普通肝素抗凝。TEG 肝素酶对比实验需同时进行 TEG 普通试验及肝素酶试验。普通试验反映患者疾病和肝素共同作用的凝血障碍;肝素酶试验破坏了血液中的肝素,仅体现患者疾病相关的凝血障碍。通过两个试验 R 时间的对比可全面准确地评估患者的凝血功能并指导肝素剂量的调整,以 R 普通检测(R_{CK})/R 肝素酶检测(R_{CKH})的比值在 1.5~2.0 为宜^[89](图 7)。如使用抗 Xa 活性来监测 SIC 患者的抗凝效果,其控制目标范围为 0.5~0.7 U/mL。




图 7 血栓弹力图肝素酶对比试验

Fig.7 Heparinase-modified thromboelastography

低分子肝素因半衰期长、不易监测,故建议在轻中度凝血功能障碍时使用。低分子肝素产品规格较多,一般建议起始剂量为 50 U/kg,每 12 h 一次静脉注射或皮下注射,并应用抗 Xa 活性监测剂量,控制目标范围为 0.3~0.5 U/mL^[90]。因为低分子肝素经肾脏代谢,肾功能不全者尤其应注意监测。当凝血功能障碍基本得到纠正(具体表现为血小板计数可自行维持在正常水平,TAT、D-二聚体等凝血指标和黏弹力试验结果基本正常)时,即可停用抗凝药物。

应用肝素类药物应尽量维持抗凝血酶活性>30%,否则可能影响抗凝效果。如果抗凝血酶活性明显降低,可通过补充新鲜冰冻血浆或抗凝血酶浓缩物提高抗凝血酶活性。应用肝素类药物还须监测血小板计数,如血小板计数明显减少甚至合并静脉、动脉血栓形成时,需考虑肝素诱导的血小板减少症(heparin-induced thrombocytopenia, HIT)。据报道,HIT 的发生率为 0.1%~5.0%,且普通肝素的诱发风险明显高于低分子肝素^[91]。HIT 的发生机制为:肝素诱导产生抗血小板第 4 因子(platelet factor 4, PF4)抗体复合物,进而激活血小板形成血小板性血栓^[92]。如发生 HIT,应立即停用肝素类药物,并改用阿加曲班或比伐卢定抗凝^[93]。现有的关于阿加曲班预防脓毒症深静脉血栓的小样本研究提示,需关注阿加曲班的药物剂量和出血并发症,但目前尚无阿加曲班或比伐卢定直接用于 SIC 抗凝治疗的证据^[94-95]。

遗憾的是,目前仍缺乏支持肝素治疗脓毒症和脓毒症性 DIC 的有效的循证医学证据。既往多项大样本随机对照研究显示,小剂量普通肝素对脓毒症的预后甚至凝血功能实验室指标并无明显影响^[96-97]。近期一项针对 MIMIC IV 数据库中 3377 例 SIC 患者的回顾性研究显

示, 普通肝素可改善 SIC 评分 ≥ 4 分的脓毒症患者的预后^[98]。应用 rhTM 治疗 SIC 的多中心随机对照试验(SCARLET 试验)结果显示, rhTM 不能改善 SIC 患者的总体预后, 但可改善 TAT 水平升高 SIC 患者的预后^[99-100]。上述研究均提示选择恰当的目标人群是抗凝治疗改善脓毒症和 SIC 预后的关键^[101]。

推荐意见 9 高出血风险 SIC 患者需抗凝时可选择甲磺酸奈莫司他(推荐强度 II, 证据等级 C)

当血管内皮严重损伤和凝血底物过度消耗时, SIC 患者可能面临高出血风险。甲磺酸奈莫司他为丝氨酸蛋白酶抑制剂, 可抑制活化的凝血因子 IIa、Xa、XIa 和组织因子 TF-VII 复合物, 同时还具有抗血小板聚集、抑制纤溶酶、抗炎和保护血管内皮的作用^[102]。甲磺酸奈莫司他在血液中可被羧酸酯酶能迅速降解, 半衰期仅 8 min, 在体内会迅速失活, 发挥抗凝作用时出血风险低, 被广泛用于体外循环和 DIC 的抗凝治疗^[103-106]。通常可通过检测患者的 APTT 或 TEG-R 时间调整甲磺酸奈莫司他的抗凝剂量。对高出血风险且行血液净化的患者, 也可在滤器后和体内同时采血, 全面评估奈莫司他的抗凝强度及出血风险^[107](图 8)。2020 年日本的一项回顾性研究显示, 奈莫司他可降低接受血液净化的脓毒症患者的病死率^[108]。此外, 作为跨膜蛋白酶丝氨酸 2(transmembrane protease serine 2, TMPRSS2)抑制剂, 奈莫司他能够抑制新型冠状病毒(severe acute respiratory syndrome coronavirus 2, SARS-CoV-2)与血管紧张素转换酶 2(angiotensin-converting enzyme 2, ACE2)结合物向细胞内的转移, 因此被认为是抑制新冠病毒传播的有效药物^[109]。已有研究显示, 奈莫司他能够改善新冠病毒感染患者的预后并减轻器官损伤^[110-111]。




图 8 高出血风险患者应用甲磺酸奈莫司他的抗凝监测

Fig.8 Anticoagulation monitoring of nafamostat mesilate in patients at high risk of bleeding

推荐意见 10 对合并高心血管血栓风险的 SIC 患者, 可根据血小板计数调整抗血小板治疗措施(推荐强度 I, 证据等级 A)

由于脓毒症早期血小板过度活化可能加重凝血紊乱和炎症失调, 因此, 有学者对脓毒症患者使用抗血小板药物能否获益开展了相关研究。多项大型回顾性队列研究显示, 接受阿司匹林治疗的全身炎症反应综合征或肺炎患者病死率较未使用者明显降低, 进展为脓毒症的发生率也明显降低, 但氯吡格雷无此效应^[112-114]。目前, 多项探讨阿司匹林能否使脓症患者获益的大型多中心随机对照研究纷纷展开。遗憾的是, 这些研究结果显示, 无论是预防或治疗, 阿司匹林对脓毒症患者的预后和器官保护均未显示出有益的作用^[115-116]。ESC 和 ISTH 共同发布的《严重感染患者抗栓治疗管理指南》推荐, 具有高心血管血栓风险的 SIC 患者需接受抗血小板药物治疗时, 需综合评估 SIC 患者的出血与血栓风险, 当血小板计数 $\leq 20 \times 10^9/L$ 时, 需停用抗血小板药物或仅维持一个抗血小板药物; 当血小板计数 $> 20 \times 10^9/L$ 时, 可考虑使用单一或双联抗血小板药物治疗^[9]。

推荐意见 11 不推荐常规使用抗纤溶药物治疗 SIC(推荐强度II, 证据等级 B)

氨甲环酸是人工合成的赖氨酸衍生物, 可竞争性结合纤溶酶原和纤溶酶的赖氨酸结合位点, 从而抑制纤维蛋白的降解, 发挥止血作用^[117]。氨甲环酸一般适用于治疗高纤溶性 DIC。脓毒症时纤溶系统功能总体呈抑制状态, 使用氨甲环酸等抗纤溶药物会加重纤溶抑制, 促进血栓形成, 导致器官功能障碍加重甚至增加死亡风险^[118]。因此, 不推荐常规使用抗纤溶药物治疗 SIC。如 SIC 患者合并大出血或原发性纤溶亢进, 需根据病情谨慎决定抗纤溶药物的使用剂量及疗程。

推荐意见 12 高出血风险或已经发生出血的 SIC 患者应进行目标导向的替代治疗(推荐强度II, 证据等级 C)

高出血风险或发生活动性出血的 SIC 患者可以凝血实验室指标为目标进行替代治疗^[119](表 6)。目前尚无 SIC 患者进行替代治疗的循证医学证据, 以下推荐意见主要依据现有的 DIC 治疗指南^[120-122]。SIC 患者进行替代治疗总体应采取个体化输血原则, 防止过早、过多输注加重血栓栓塞。

表 6 高出血风险或活动性出血的 SIC 患者的替代治疗

Tab.6 Alternative treatments for SIC patients with high bleeding risk or active bleeding

治疗目的	启动指征	替代措施
补充凝血因子	PT 或 APTT 延长>1.5 倍; 或 TEG-R>15 min; 或凝血与血小板功能分析 ACT>240 s	新鲜冰冻血浆(10~15 mL/kg)或 4 种凝血因子凝血酶原复合物(10~15 U/kg)
补充纤维蛋白原	纤维蛋白原<1.0 g/L; 或 TEG 功能纤维蛋白原检测指标 FFMA<10 mm; 或凝血与血小板功能分析 CR<10	冷沉淀(10 mL/kg)或人纤维蛋白原(30~50 mg/kg)
	非出血的 SIC 患者血小板计数<20×10 ⁹ /L	
	发生活动性出血且血小板计数<50×10 ⁹ /L	
补充血小板	正在进行 ECMO 治疗且血小板计数<80×10 ⁹ /L TEG-MA<43 mm 且纤维蛋白原>1 g/L 凝血与血小板功能分析 PF<1	机采血小板 1~2 个治疗量

注: SIC. 脓毒症性凝血病; TEG. 血栓弹力图; APTT. 活化部分凝血活酶时间; PT. 凝血酶原时间; FFMA. 功能性纤维蛋白原试验的最大振幅; CR. 凝血速率; ECMO. 体外膜氧合肺; PF. 血小板指数

推荐意见 13 如 SIC 发展成脓毒症性 DIC, 可行血浆置换(推荐强度II, 证据等级 C)

血浆置换是先将患者血液引至体外, 经离心法或膜分离法分离血浆和细胞成分, 然后弃去血浆, 再把细胞成分及所需补充的新鲜血浆回输体内^[123]。血浆置换既能清除大量炎性介质, 又能去除患者血浆中的胆红素等毒性物质, 同时补充凝血因子以改善危重患者的凝血状态, 是治疗脓毒症性 DIC 的有效方法^[124]。一项针对 102 名欧洲重症医师的网络调查显示, 34.1%的医师认为血浆置换是治疗脓毒症休克最常用的方法, 且 51.5%的医师认为其有效^[125]。多项纳入随机对照研究和队列研究的荟萃分析显示, 接受血浆置换治疗的脓毒症合并严重器官功能障碍患者的短期生存率明显高于标准治疗组^[126-127], 其机制主要与改善血管内皮细胞功能有关^[128]。此外, 一项纳入 851 例新冠重症患者的回顾性研究显示, 入院 96 h 内给予甲泼尼龙、机械通气和血浆置换“三联”疗法可改善患者预后^[129]。血浆置换的剂量通常以每次 1.2~1.5 倍血浆量为宜, 具体频次应根据患者对血浆置换的治疗反应和器官功能损害的严

重程度而定。

推荐意见 14 治疗 SIC 时应预防稀释性凝血病(推荐强度II, 证据等级 C)

液体复苏是治疗脓毒症的基本技术。指南明确推荐,对于脓毒症导致的低血压或脓毒症休克患者,在最初 3 h 内应静脉输注至少 30 mL/kg 的晶体液^[51]。但对于初始液体复苏后仍有低灌注和容量不足的脓毒症休克患者,尚无证据证实后续 24 h 内应采用限制性还是开放性液体管理策略^[51,130]。已有研究证实,脓毒症时高容量液体复苏会导致血管内皮功能损害、凝血时间延长和器官功能障碍,甚至增高病死率^[131-134]。因此, SIC 患者进行液体复苏时应加强液体管理,避免发生稀释性凝血病^[135-136]。

6 展望

本共识依据最新研究进展和专家诊疗经验,从发病机制上介绍了免疫血栓与血栓炎症的序贯过程,从分型上提出了高凝血症与低凝血症,从诊断上提出了适合中国人群的 SIC 诊断方法,从治疗上提出了“三抗一防”的治疗策略。但因为 SIC 的机制复杂,异质性大,同时缺乏有力的循证医学依据,本共识仍存在以下不足:(1)对免疫血栓和血栓炎症的认识仍停留在理论阶段,临床上尚缺乏可靠的识别方法;(2)虽然介绍了脓毒症性高凝血症和低凝血症的典型实验室表现,但因为缺乏循证依据,没有推荐明确的诊断标准;(3)目前的 SIC 诊断方法尚未包括内皮细胞损伤分子标志物,还需要持续优化并进行临床验证;(4)SIC 伴出血时的替代治疗方案的具体标准仍需要开展临床研究进行验证。

编写组成员:

顾问(按拼音排序):李维勤(解放军东部战区总医院重症医学科);林洪远(解放军总医院第四医学中心重症医学科);马晓春(中国医科大学第一附属医院重症医学科);宋青(解放军总医院海南医院重症医学科);杨仁池(中国医学科学院血液病医院血栓与止血中心)

编委会成员(按拼音排序):陈淼(海南医学院第一附属医院急诊科);程鹏(广西医科大学第一附属医院血液科);戴菁(上海交通大学附属瑞金医院检验科);刁孟元(杭州市第一人民医院重症医学科);丁仁彧(中国医科大学第一附属医院重症医学科);房云海(山东省血液中心血友病诊疗中心);桂培根(湖南师范大学附属长沙医院急重症医学部);郭军(四川大学华西医院重症医学科);贾宝辉(郑州大学附属郑州市中心医院重症医学科);柯路(解放军东部战区总医院重症医学科);赖冬(厦门市第二医院输血科);李传保(北京医院检验科);吕奔(中南大学附属湘雅二医院);马林浩(海军军医大学附属长征医院急救科);梅恒(华中科技大学医学院附属协和医院血液科);潘景业(温州医科大学附属第一医院重症医学科);施贤清(贵州省人民医院重症医学科);寿松涛(天津医科大学附属天津总医院急诊科);宋景春(解放军联勤保障部队第 908 医院重症医学科);宋振举(复旦大学附属中山医院急诊科);唐宁(华中科技大学医学院附属同济医院检验科);王岗(西安交通大学第二附属医院重症医学科);王昊(山东大学附属齐鲁医院重症医学科);王秋实(中国医科大学附属盛京医院输血科);吴俊(首都医科大学附属北京积水潭医院检验科);杨军(武汉亚洲心脏病医院检验科);尹海燕(暨南大学附属第一医院重症医学科);张根生(浙江大学医学院第二附属医院重症医学科);张进华(福建省妇幼保健院药学科);张磊(西安交通大学第二附属医院检验科);张伟(解放军联勤保障部队第 900 医院急诊科);周静(四川大学华西医院检验科);朱峰(同济大学附属东方医院重症医学科)

秘书:钟林翠(解放军联勤保障部队第 908 医院重症医学科);张小玲(西安交通大学第二附属医院重症医学科)

【参考文献】

- [1] Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3)[J]. *JAMA*, 2016, 315(8): 801-810.
- [2] Iba T, Nisio MD, Levy JH, et al. New criteria for sepsis-induced coagulopathy (SIC) following the revised sepsis definition: a retrospective analysis of a nationwide survey[J]. *BMJ Open*, 2017, 7(9): e017046.
- [3] Schmoch T, Möhnlé P, Weigand MA, et al. The prevalence of sepsis-induced coagulopathy in patients with sepsis - a secondary analysis of two German multicenter randomized controlled trials[J]. *Ann Intensive Care*, 2023, 13(1): 3.
- [4] Yamakawa K, Yoshimura J, Ito T, et al. External validation of the two newly proposed criteria for assessing coagulopathy in sepsis[J]. *Thromb Haemost*, 2019, 119(2): 203-212.
- [5] Tanaka C, Tagami T, Kudo S, et al. Validation of sepsis-induced coagulopathy score in critically ill patients with septic shock: post hoc analysis of a nationwide multicenter observational study in Japan[J]. *Int J Hematol*, 2021, 114(2): 164-171.
- [6] Singh B, Hanson AC, Alhurani R, et al. Trends in the incidence and outcomes of disseminated intravascular coagulation in critically ill patients (2004-2010): a population-based study[J]. *Chest*, 2013, 143(5): 1235-1242.
- [7] Ding R, Wang Z, Lin Y, et al. Comparison of a new criteria for sepsis-induced coagulopathy and International Society on Thrombosis and Haemostasis disseminated intravascular coagulation score in critically ill patients with sepsis 3.0: a retrospective study[J]. *Blood Coagul Fibrinolysis*, 2018, 29(6): 551-558.
- [8] Iba T, Levi M, Thachil J, et al. Communication from the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis on sepsis-induced coagulopathy in the management of sepsis[J]. *J Thromb Haemost*, 2023, 21(1): 145-153.
- [9] Gigante B, Levy JH, van Gorp E, et al. Management of patients on antithrombotic therapy with severe infections: a joint clinical consensus statement of the ESC Working Group on Thrombosis, the ESC Working Group on Atherosclerosis and Vascular Biology, and the International Society on Thrombosis and Haemostasis[J]. *Eur Heart J*, 2023, 44(32): 3040-3058.
- [10] Zaid Y, Merhi Y. Implication of platelets in immuno-thrombosis and thrombo-inflammation[J]. *Front Cardiovasc Med*, 2022, 9: 863846.
- [11] Engelmann B, Massberg S. Thrombosis as an intravascular effector of innate immunity[J]. *Nat Rev Immunol*, 2013, 13(1): 34-45.
- [12] Swystun LL, Liaw PC. The role of leukocytes in thrombosis[J]. *Blood*, 2016, 128(6): 753-762.
- [13] Iba T, Levi M, Levy JH. Intracellular communication and immunothrombosis in sepsis[J]. *J Thromb Haemost*, 2022, 20(11): 2475-2484.
- [14] Iba T, Levi M, Thachil J, et al. Disseminated intravascular coagulation: the past, present, and future considerations[J]. *Semin Thromb Hemost*, 2022, 48(8): 978-987.
- [15] Yang S, Qi H, Kan K, et al. Neutrophil extracellular traps promote hypercoagulability in patients with sepsis[J]. *Shock*, 2017, 47(2): 132-139.
- [16] Wada T, Gando S. Phenotypes of disseminated intravascular coagulation[J]. *Thromb Haemost*, 2024, 124(3): 181-191.
- [17] Musgrave KM, Scott J, Sendama W, et al. Tissue factor expression in monocyte subsets during human immunothrombosis, endotoxemia and sepsis[J]. *Thromb Res*, 2023, 228: 10-20.

- [18] Ryan TAJ, Preston RJS, O'Neill LAJ. Immunothrombosis and the molecular control of tissue factor by pyroptosis: prospects for new anticoagulants[J]. *Biochem J*, 2022, 479(6): 731-750.
- [19] Shi J, Tang Y, Liang F, et al. NLRP3 inflammasome contributes to endotoxin-induced coagulation[J]. *Thromb Res*, 2022, 214: 8-15.
- [20] Iba T, Umemura Y, Wada H, et al. Roles of coagulation abnormalities and microthrombosis in sepsis: pathophysiology, diagnosis, and treatment[J]. *Arch Med Res*, 2021, 52(8): 788-797.
- [21] Masuda T, Shoko T. Clinical investigation of the utility of a pair of coagulation-fibrinolysis markers for definite diagnosis of sepsis-induced disseminated intravascular coagulation: a single-center, diagnostic, prospective, observational study[J]. *Thromb Res*, 2020, 192: 116-121.
- [22] Ninan K F, Iyadurai R, Varghese JK, et al. Thromboelastograph: a prognostic marker in sepsis with organ dysfunction without overt bleeding[J]. *J Crit Care*, 2021, 65: 177-183.
- [23] Premkumar M, Loganathan S, Kajal K, et al. COVID-19-related dynamic coagulation disturbances and anticoagulation strategies using conventional D-dimer and point-of-care Sonoclot tests: a prospective cohort study[J]. *BMJ Open*, 2022, 12(5): e051971.
- [24] Lipets EN, Ataullakhanov FI. Global assays of hemostasis in the diagnostics of hypercoagulation and evaluation of thrombosis risk[J]. *Thromb J*, 2015, 13(1): 4.
- [25] Zhong L, Dou J, Lin Q, et al. Tissue-type plasminogen activator-inhibitor complex as an early predictor of septic shock: a retrospective, single-center study[J]. *Dis Markers*, 2022, 2022: 9364037.
- [26] Boscolo A, Spiezia L, De Cassai A, et al. Are thromboelastometric and thromboelastographic parameters associated with mortality in septic patients? A systematic review and meta-analysis[J]. *J Crit Care*, 2021, 61: 5-13.
- [27] Kim SM, Kim SI, Yu G, et al. Role of thromboelastography as an early predictor of disseminated intravascular coagulation in patients with septic shock[J]. *J Clin Med*, 2020, 9(12): 3883.
- [28] Taylor FB Jr, Toh CH, Hoots WK, et al. Towards definition, clinical and laboratory criteria, and a scoring system for disseminated intravascular coagulation[J]. *Thromb Haemost*, 2001, 86(5): 1327-1330.
- [29] Meziani F, Iba T, Levy JH, et al. Sepsis-induced coagulopathy: a matter of timeline[J]. *Intensive Care Med*, 2024, 50(8): 1404-1405.
- [30] Thachil J, Iba T. Designing the diagnostic criteria for disseminated intravascular coagulation (DIC)[J]. *Juntendo Iji Zasshi*, 2023, 69(6): 463-465.
- [31] Iba T, Levy JH, Warkentin TE, et al. Diagnosis and management of sepsis-induced coagulopathy and disseminated intravascular coagulation[J]. *J Thromb Haemost*, 2019, 17(11): 1989-1994.
- [32] 中国医药教育协会. T/CMEAS 019-2024. 凝血障碍诊断规范[S]. 北京: 中国标准出版社, 2024, 1-10.
- [33] Iba T, Arakawa M, Di Nisio M, et al. Newly proposed sepsis-induced coagulopathy precedes international society on thrombosis and haemostasis overt-disseminated intravascular coagulation and predicts high mortality[J]. *J Intensive Care Med*, 2020, 35(7): 643-649.
- [34] 钟林翠, 宋景春, 吴骏, 等. 脓毒症性凝血病修正评分系统的临床价值分析[J]. *血栓与止血学*, 2024, 30(2): 52-57.
- [35] Rhodes A, Evans LE, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016[J]. *Intensive Care Med*, 2017, 43(3): 304-377.
- [36] Seymour CW, Gesten F, Prescott HC, et al. Time to treatment and mortality during mandated emergency care for sepsis[J]. *N Engl J Med*, 2017, 376(23): 2235-2244.
- [37] Martínez ML, Ferrer R, Torrents E, et al. Impact of source control in patients with severe sepsis

- and septic shock[J]. *Crit Care Med*, 2017, 45(1): 11-19.
- [38] De Waele JJ. Importance of timely and adequate source control in sepsis and septic shock[J]. *J Intensive Med*, 2024, 4(3): 281-286.
- [39] Lester W, Bent C, Alikhan R, et al. A British Society for Haematology guideline on the assessment and management of bleeding risk prior to invasive procedures[J]. *Br J Haematol*, 2024, 204(5): 1697-1713.
- [40] Wang Q, Liang P, Xu Y, et al. Serum trough concentration threshold and risk factors of cefoperazone-induced coagulopathy in critically ill patients: a retrospective case-control study[J]. *Eur J Clin Pharmacol*, 2024, 80(5): 737-746.
- [41] Yu Z, Chen H. Piperacillin/tazobactam-induced coagulopathy in a patient through a vitamin K-dependent mechanism[J]. *Eur J Hosp Pharm*, 2021, 28(4): 237-238.
- [42] Tiperneni R, Khalid F, Fichadiya H, et al. Deranged haemostasis: rifampin-induced coagulopathy[J]. *Eur J Case Rep Intern Med*, 2022, 9(5): 003380.
- [43] Wang D, Lin C, Gu C, et al. Tigecycline-associated coagulopathy: a single-center retrospective analysis[J]. *Pharmacology*, 2022, 107(9-10): 524-536.
- [44] Al Qamariat Z, Aljaffar AA, Alabdulaal ZS, et al. Rapid onset and recovery linezolid-induced thrombocytopenia: a large-sample, single-center retrospective cohort study[J]. *Drug Healthc Patient Saf*, 2024, 16: 43-49.
- [45] Mitta A, Curtis BR, Reese JA, et al. Drug-induced thrombocytopenia: 2019 Update of clinical and laboratory data[J]. *Am J Hematol*, 2019, 94(3): E76-E78.
- [46] Vayne C, Guéry EA, Rollin J, et al. Pathophysiology and diagnosis of drug-induced immune thrombocytopenia[J]. *J Clin Med*, 2020, 9(7): 2212.
- [47] Nedeva C, Menassa J, Puthalakath H. Sepsis: inflammation is a necessary evil[J]. *Front Cell Dev Biol*, 2019, 7: 108.
- [48] Bennett IL Jr, Finland M, Hamburger M, et al. A double-blind study of the effectiveness of cortisol in the management of severe infections[J]. *Trans Assoc Am Physicians*, 1962, 75: 198-207.
- [49] Rochwerg B, Oczkowski SJ, Siemieniuk RAC, et al. Corticosteroids in sepsis: an updated systematic review and meta-analysis[J]. *Crit Care Med*, 2018, 46(9): 1411-1420.
- [50] Venkatesh B, Finfer S, Cohen J, et al. Adjunctive glucocorticoid therapy in patients with septic shock[J]. *N Engl J Med*, 2018, 378(9): 797-808.
- [51] Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021[J]. *Intensive Care Med*, 2021, 47(11): 1181-1247.
- [52] Chaudhuri D, Nei AM, Rochwerg B, et al. 2024 focused update: guidelines on use of corticosteroids in sepsis, acute respiratory distress syndrome, and community-acquired pneumonia[J]. *Crit Care Med*, 2024, 52(5): e219-e233.
- [53] Schmidt C, Weißmüller S, Bohländer F, et al. The dual role of a polyvalent IgM/IgA-enriched immunoglobulin preparation in activating and inhibiting the complement system[J]. *Biomedicines*, 2021, 9(7): 817.
- [54] Asakura H, Takahashi Y, Kubo A, et al. Immunoglobulin preparations attenuate organ dysfunction and hemostatic abnormality by suppressing the production of cytokines in lipopolysaccharide-induced disseminated intravascular coagulation in rats[J]. *Crit Care Med*, 2006, 34(9): 2421-2425.
- [55] Ishikura H, Nakamura Y, Kawano Y, et al. Intravenous immunoglobulin improves sepsis-

- induced coagulopathy: a retrospective, single-center observational study[J]. *J Crit Care*, 2015, 30(3): 579-583.
- [56] Wand S, Klages M, Kirbach C, et al. IgM-enriched immunoglobulin attenuates systemic endotoxin activity in early severe sepsis: a before-after cohort study[J]. *PLoS One*, 2016, 11(8): e0160907.
- [57] Busani S, Damiani E, Cavazzuti I, et al. Intravenous immunoglobulin in septic shock: review of the mechanisms of action and meta-analysis of the clinical effectiveness[J]. *Minerva Anesthesiol*, 2016, 82(5): 559-572.
- [58] Cui J, Wei X, Lv H, et al. The clinical efficacy of intravenous IgM-enriched immunoglobulin (pentaglobin) in sepsis or septic shock: a meta-analysis with trial sequential analysis[J]. *Ann Intensive Care*, 2019, 9(1): 27.
- [59] Schmidt C, Weißmüller S, Heinz CC. Multifaceted tissue-protective functions of polyvalent immunoglobulin preparations in severe infections-interactions with neutrophils, complement, and coagulation pathways[J]. *Biomedicines*, 2023, 11(11): 3022.
- [60] Mandl J, Szarka A, Bánhegyi G. Vitamin C: update on physiology and pharmacology[J]. *Br J Pharmacol*, 2009, 157(7): 1097-1110.
- [61] Oudemans-van Straaten HM, Spoelstra-de Man AM, de Waard MC. Vitamin C revisited[J]. *Crit Care*, 2014, 18(4): 460.
- [62] Kuhn SO, Meissner K, Mayes LM, et al. Vitamin C in sepsis[J]. *Curr Opin Anaesthesiol*, 2018, 31(1): 55-60.
- [63] Borrelli E, Roux-Lombard P, Grau GE, et al. Plasma concentrations of cytokines, their soluble receptors, and antioxidant vitamins can predict the development of multiple organ failure in patients at risk[J]. *Crit Care Med*, 1996, 24(3): 392-397.
- [64] Marik PE, Khangoora V, Rivera R, et al. Hydrocortisone, vitamin C, and thiamine for the treatment of severe sepsis and septic shock: a retrospective before-after study[J]. *Chest*, 2017, 151(6): 1229-1238.
- [65] Putzu A, Daems AM, Lopez-Delgado JC, et al. The effect of vitamin C on clinical outcome in critically ill patients: a systematic review with meta-analysis of randomized controlled trials[J]. *Crit Care Med*, 2019, 47(6): 774-783.
- [66] Martimbianco ALC, Pacheco RL, Bagattini ÂM, et al. Vitamin C-based regimens for sepsis and septic shock: systematic review and meta-analysis of randomized clinical trials[J]. *J Crit Care*, 2022, 71: 154099.
- [67] Fujii T, Luethi N, Young PJ, et al. Effect of vitamin C, hydrocortisone, and thiamine vs hydrocortisone alone on time alive and free of vasopressor support among patients with septic shock: the VITAMINS randomized clinical trial[J]. *JAMA*, 2020, 323(5): 423-431.
- [68] Moskowitz A, Huang DT, Hou PC, et al. Effect of ascorbic acid, corticosteroids, and thiamine on organ injury in septic shock: the ACTS randomized clinical trial[J]. *JAMA*, 2020, 324(7): 642-650.
- [69] Linder A, Russell JA. An exciting candidate therapy for sepsis: ulinastatin, a urinary protease inhibitor[J]. *Intensive Care Med*, 2014, 40(8): 1164-1167.
- [70] Shu H, Liu K, He Q, et al. Ulinastatin, a protease inhibitor, may inhibit allogeneic blood transfusion-associated pro-inflammatory cytokines and systemic inflammatory response syndrome and improve postoperative recovery[J]. *Blood Transfus*, 2014, 12(Suppl 1): s109-s118.

- [71] Inoue KI, Takano H. Urinary trypsin inhibitor as a therapeutic option for endotoxin-related inflammatory disorders[J]. *Expert Opin Investig Drugs*, 2010, 19(4): 513-520.
- [72] Wang H, Liu B, Tang Y, et al. Improvement of sepsis prognosis by ulinastatin: a systematic review and Meta-analysis of randomized controlled trials[J]. *Front Pharmacol*, 2019, 10: 1370.
- [73] Zhang J, Tian J, Sun H, et al. How does continuous renal replacement therapy affect septic acute kidney injury?[J]. *Blood Purif*, 2018, 46(4): 326-331.
- [74] Murugan R, Wen X, Shah N, et al. Plasma inflammatory and apoptosis markers are associated with dialysis dependence and death among critically ill patients receiving renal replacement therapy[J]. *Nephrol Dial Transplant*, 2014, 29(10): 1854-1864.
- [75] Atan R, Crosbie DC, Bellomo R. Techniques of extracorporeal cytokine removal: a systematic review of human studies[J]. *Ren Fail*, 2013, 35(8): 1061-1070.
- [76] Lumlertgul N, Hall A, Camporota L, et al. Clearance of inflammatory cytokines in patients with septic acute kidney injury during renal replacement therapy using the EMiC2 filter (Clic-AKI study)[J]. *Crit Care*, 2021, 25(1): 39.
- [77] Onichimowski D, Goraj R, Jalali R, et al. Practical issues of nutrition during continuous renal replacement therapy[J]. *Anaesthesiol Intensive Ther*, 2017, 49(4): 309-316.
- [78] Levi M, van der Poll T. Coagulation in patients with severe sepsis[J]. *Semin Thromb Hemost*, 2015, 41(1): 9-15.
- [79] Wu S, Xu T, Wu C, et al. Continuous renal replacement therapy in sepsis-associated acute kidney injury: effects on inflammatory mediators and coagulation function[J]. *Asian J Surg*, 2021, 44(10): 1254-1259.
- [80] Zou F, Tang X, Lei X, et al. Treatment efficacy of continuous renal replacement on symptoms, inflammatory mediators, and coagulation function in patients with sepsis-associated acute kidney injury[J]. *Arch Esp Urol*, 2022, 75(9): 746-752.
- [81] Cruz DN, Antonelli M, Fumagalli R, et al. Early use of polymyxin B hemoperfusion in abdominal septic shock: the EUPHAS randomized controlled trial[J]. *JAMA*, 2009, 301(23): 2445-2452.
- [82] Osawa I, Goto T, Kudo D, et al. Targeted therapy using polymyxin B hemadsorption in patients with sepsis: a post-hoc analysis of the JSEPTIC-DIC study and the EUPHRATES trial[J]. *Crit Care*, 2023, 27(1): 245.
- [83] Klein DJ, Foster D, Walker PM, et al. Polymyxin B hemoperfusion in endotoxemic septic shock patients without extreme endotoxemia: a post hoc analysis of the EUPHRATES trial[J]. *Intensive Care Med*, 2018, 44(12): 2205-2212.
- [84] Dellinger RP, Bagshaw SM, Antonelli M, et al. Effect of targeted polymyxin B hemoperfusion on 28-day mortality in patients with septic shock and elevated endotoxin level: the EUPHRATES randomized clinical trial[J]. *JAMA*, 2018, 320(14): 1455-1463.
- [85] Umemura Y, Yamakawa K. Optimal patient selection for anticoagulant therapy in sepsis: an evidence-based proposal from Japan[J]. *J Thromb Haemost*, 2018, 16(3): 462-464.
- [86] Yamakawa K, Umemura Y, Hayakawa M, et al. Benefit profile of anticoagulant therapy in sepsis: a nationwide multicentre registry in Japan[J]. *Crit Care*, 2016, 20(1): 229.
- [87] Murao A, Kato T, Yamane T, et al. Benefit profile of thrombomodulin Alfa combined with antithrombin concentrate in patients with sepsis-induced disseminated intravascular coagulation[J]. *Clin Appl Thromb Hemost*, 2022, 28: 10760296221077096.
- [88] 中华医学会血液学分会血栓与止血学组. 弥散性血管内凝血诊断与治疗中国专家共识

- (2012 年版)[J]. 中华血液学杂志, 2012, 33(11): 978-979.
- [89] Coppell JA, Thalheimer U, Zambruni A, et al. The effects of unfractionated heparin, low molecular weight heparin and danaparoid on the thromboelastogram (TEG): an *in-vitro* comparison of standard and heparinase-modified TEGs with conventional coagulation assays[J]. Blood Coagul Fibrinolysis, 2006, 17(2): 97-104.
- [90] Samimi MN, Hale A, Schults J, et al. Clinical guidance for unfractionated heparin dosing and monitoring in critically ill patients[J]. Expert Opin Pharmacother, 2024, 25(8): 985-997.
- [91] Cuker A, Arepally GM, Chong BH, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: heparin-induced thrombocytopenia[J]. Blood Adv, 2018, 2(22): 3360-3392.
- [92] Alhanshani AA. Heparin induced thrombocytopenia - pathophysiology, diagnosis and treatment: a narrative review[J]. Int J Gen Med, 2023, 16: 3947-3953.
- [93] Greinacher A. CLINICAL PRACTICE. Heparin-induced thrombocytopenia[J]. N Engl J Med, 2015, 373(3): 252-261.
- [94] Bachler M, Asmis LM, Koscielny J, et al. Thromboprophylaxis with argatroban in critically ill patients with sepsis: a review[J]. Blood Coagul Fibrinolysis, 2022, 33(5): 239-256.
- [95] Heubner L, Oertel R, Tiebel O, et al. Monitoring of argatroban in critically ill patients: a prospective study comparing activated partial thromboplastin time, point-of-care viscoelastic testing with ecarin clotting time and diluted thrombin time to mass spectrometry[J]. Anesthesiology, 2024, 140(2): 261-271.
- [96] Polderman KH, Girbes AR. Drug intervention trials in sepsis: divergent results[J]. Lancet, 2004, 363(9422): 1721-1723.
- [97] Jaimes F, de la Rosa G, Morales C, et al. Unfractionated heparin for treatment of sepsis: a randomized clinical trial (The HETRASE Study)[J]. Crit Care Med, 2009, 37(4): 1185-1196.
- [98] Huang JJ, Zou ZY, Zhou ZP, et al. Effectiveness of early heparin therapy on outcomes in critically ill patients with sepsis-induced coagulopathy[J]. Front Pharmacol, 2023, 14: 1173893.
- [99] Vincent JL, Francois B, Zabolotskikh I, et al. Effect of a recombinant human soluble thrombomodulin on mortality in patients with sepsis-associated coagulopathy: the SCARLET randomized clinical trial[J]. JAMA, 2019, 321(20): 1993-2002.
- [100] Levi M, Vincent JL, Tanaka K, et al. Effect of a recombinant human soluble thrombomodulin on baseline coagulation biomarker levels and mortality outcome in patients with sepsis-associated coagulopathy[J]. Crit Care Med, 2020, 48(8): 1140-1147.
- [101] Umemura Y, Yamakawa K, Ogura H, et al. Efficacy and safety of anticoagulant therapy in three specific populations with sepsis: a meta-analysis of randomized controlled trials[J]. J Thromb Haemost, 2016, 14(3): 518-530.
- [102] Quinn TM, Gaughan EE, Bruce A, et al. Randomised controlled trial of intravenous nafamostat mesylate in COVID pneumonitis: phase 1b/2a experimental study to investigate safety, Pharmacokinetics and Pharmacodynamics[J]. EBioMedicine, 2022, 76: 103856.
- [103] Takahashi W, Yoneda T, Koba H, et al. Potential mechanisms of nafamostat therapy for severe COVID-19 pneumonia with disseminated intravascular coagulation[J]. Int J Infect Dis, 2021, 102: 529-531.
- [104] Minakata D, Fujiwara SI, Hayakawa J, et al. Comparison of danaparoid sodium and synthetic protease inhibitors for the treatment of disseminated intravascular coagulation associated with hematological malignancies: a retrospective analysis[J]. Acta Haematol, 2020, 143(3): 250-

- [105] Minakata D, Fujiwara SI, Ikeda T, et al. Comparison of gabexate mesilate and nafamostat mesilate for disseminated intravascular coagulation associated with hematological malignancies[J]. *Int J Hematol*, 2019, 109(2): 141-146.
- [106] Kusuzawa K, Suzuki K, Okada H, et al. Measuring the concentration of serum syndecan-1 to assess vascular endothelial glycocalyx injury during hemodialysis[J]. *Front Med*, 2021, 8: 791309.
- [107] 林青伟, 宋景春, 彭恩兰, 等. 应用甲磺酸萘莫司他治疗脓毒症性凝血病的探讨[J]. *血栓与止血学*, 2023, 29(3): 138-142.
- [108] Kamijo H, Mochizuki K, Nakamura Y, et al. Nafamostat mesylate improved survival outcomes of sepsis patients who underwent blood purification: a nationwide registry study in Japan[J]. *J Clin Med*, 2020, 9(8): 2629.
- [109] Hernández-Mitre MP, Tong SYC, Denholm JT, et al. Nafamostat mesylate for treatment of COVID-19 in hospitalised patients: a structured, narrative review[J]. *Clin Pharmacokinet*, 2022, 61(10): 1331-1343.
- [110] Morpeth SC, Venkatesh B, Totterdell JA, et al. A randomized trial of nafamostat for covid-19[J]. *NEJM Evid*, 2023, 2(11): EVIDoa2300132.
- [111] Okugawa S, Ikeda M, Kashiwabara K, et al. Antiviral effect and safety of nafamostat mesilate in patients with mild early-onset COVID-19: an exploratory multicentre randomized controlled clinical trial[J]. *Int J Antimicrob Agents*, 2023, 62(3): 106922.
- [112] Erlich JM, Talmor DS, Cartin-Ceba R, et al. Prehospitalization antiplatelet therapy is associated with a reduced incidence of acute lung injury: a population-based cohort study[J]. *Chest*, 2011, 139(2): 289-295.
- [113] Eisen DP, Reid D, McBryde ES. Acetyl salicylic acid usage and mortality in critically ill patients with the systemic inflammatory response syndrome and sepsis[J]. *Crit Care Med*, 2012, 40(6): 1761-1767.
- [114] Falcone M, Russo A, Cangemi R, et al. Lower mortality rate in elderly patients with community-onset pneumonia on treatment with aspirin[J]. *J Am Heart Assoc*, 2015, 4(1): e001595.
- [115] Eisen DP, Leder K, Woods RL, et al. Effect of aspirin on deaths associated with sepsis in healthy older people (ANTISEPSIS): a randomised, double-blind, placebo-controlled primary prevention trial[J]. *Lancet Respir Med*, 2021, 9(2): 186-195.
- [116] REMAP-CAP Writing Committee for the REMAP-CAP Investigators, Bradbury CA, Lawler PR, et al. Effect of antiplatelet therapy on survival and organ support-free days in critically ill patients with COVID-19: a randomized clinical trial[J]. *JAMA*, 2022, 327(13): 1247-1259.
- [117] Patel PA, Wyrobek JA, Butwick AJ, et al. Update on applications and limitations of perioperative tranexamic acid[J]. *Anesth Analg*, 2022, 135(3): 460-473.
- [118] Napolitano F, Giudice V, Selleri C, et al. Plasminogen system in the pathophysiology of sepsis: upcoming biomarkers[J]. *Int J Mol Sci*, 2023, 24(15): 12376.
- [119] Iba T. Harmonized guidance for disseminated intravascular coagulation from the International Society on Thrombosis and Haemostasis and the current status of anticoagulant therapy in Japan[J]. *J Thromb Haemost*, 2013, 11(11): 2076-2078.
- [120] Di Nisio M, Baudo F, Cosmi B, et al. Diagnosis and treatment of disseminated intravascular coagulation: guidelines of the Italian Society for Haemostasis and Thrombosis (SISET)[J].

Thromb Res, 2012, 129(5): e177-e184.

- [121] Levi M, Toh CH, Thachil J, et al. Guidelines for the diagnosis and management of disseminated intravascular coagulation. British Committee for Standards in Haematology[J]. Br J Haematol, 2009, 145(1): 24-33.
- [122] Iba T, Asakura H. Comparison between British and Japanese guidelines for the diagnosis and treatment of disseminated intravascular coagulation[J]. Br J Haematol, 2010, 149(3): 461-462.
- [123] Schwartz J, Padmanabhan A, Aqui N, et al. Guidelines on the use of therapeutic apheresis in clinical practice-evidence-based approach from the Writing Committee of the American Society for Apheresis: the seventh special issue[J]. J Clin Apher, 2016, 31(3):149-162.
- [124] Aydin K, Korkmaz S, Erkurt MA, et al. Apheresis in patients with sepsis: a multicenter retrospective study[J]. Transfus Apher Sci, 2021, 60(5): 103239.
- [125] Stahl K, Bode C, Seeliger B, et al. Current clinical practice in using adjunctive extracorporeal blood purification in sepsis and septic shock: results from the ESICM “EXPLORATION” survey[J]. Intensive Care Med Exp, 2024, 12(1): 5.
- [126] Kuklin V, Sovershaev M, Bjerner J, et al. Influence of therapeutic plasma exchange treatment on short-term mortality of critically ill adult patients with sepsis-induced organ dysfunction: a systematic review and meta-analysis[J]. Crit Care, 2024, 28(1): 12.
- [127] Lee OPE, Kanesan N, Leow EH, et al. Survival benefits of therapeutic plasma exchange in severe sepsis and septic shock: a systematic review and meta-analysis[J]. J Intensive Care Med, 2023, 38(7): 598-611.
- [128] Weng JT, Chen M, Fang DX, et al. Therapeutic plasma exchange protects patients with sepsis-associated disseminated intravascular coagulation by improving endothelial function[J]. Clin Appl Thromb Hemost, 2021, 27: 10760296211053313.
- [129] Keith P, Bohn RIC, Nguyen T, et al. Improved survival in COVID-19 related sepsis and ARDS treated with a unique “triple therapy” including therapeutic plasma exchange: a single center retrospective analysis[J]. J Clin Apher, 2024, 39(1): e22107.
- [130] Marik PE, Linde-Zwirble WT, Bittner EA, et al. Fluid administration in severe sepsis and septic shock, patterns and outcomes: an analysis of a large national database[J]. Intensive Care Med, 2017, 43(5): 625-632.
- [131] Boyd JH, Forbes J, Nakada TA, et al. Fluid resuscitation in septic shock: a positive fluid balance and elevated central venous pressure are associated with increased mortality[J]. Crit Care Med, 2011, 39(2): 259-265.
- [132] Alphonsus CS, Rodseth RN. The endothelial glycocalyx: a review of the vascular barrier[J]. Anaesthesia, 2014, 69(7): 777-784.
- [133] Zhao H, Cai X, Liu N, et al. Thromboelastography as a tool for monitoring blood coagulation dysfunction after adequate fluid resuscitation can predict poor outcomes in patients with septic shock[J]. J Chin Med Assoc, 2020, 83(7): 674-677.
- [134] Huang ACC, Lee TYT, Ko MC, et al. Fluid balance correlates with clinical course of multiple organ dysfunction syndrome and mortality in patients with septic shock[J]. PLoS One, 2019, 14(12): e0225423.
- [135] Govero AB, Yarrarapu SNS, Harrison MF, et al. Surviving sepsis guideline-directed fluid resuscitation: an assessment of practice patterns and impact on patient outcomes[J]. Crit Care Explor, 2022, 4(7): e0739.
- [136] Arabi YM, Belley-Cote E, Carsetti A, et al. European Society of Intensive Care Medicine

clinical practice guideline on fluid therapy in adult critically ill patients. Part 1: the choice of resuscitation fluids[J]. Intensive Care Med, 2024, 50(6): 813-831.

(责任编辑: 张小利)

